

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representation of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## **IMAGES ARE BEST AVAILABLE COPY**

**As rescanning documents *will not* correct images, please do not report the images to the Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

CI

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 466 222 A1**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 91201436.2

(51) Int. Cl.<sup>5</sup>: **C07K 15/10, C12N 15/29,  
C12N 5/10**

(22) Date of filing: 10.06.91

(30) Priority: 11.06.90 US 535636

(43) Date of publication of application:  
15.01.92 Bulletin 92/03

(94) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IT LI LU NL SE**

(71) Applicant: **DOWELANCO**  
9002 Purdue Road  
Indianapolis, Indiana 46268-3030(US)

(72) Inventor: **Walsh, Terence A.**  
E. Old Pine Trail  
Midland, Michigan 48640(US)  
Inventor: **Hey, Timothy D.**  
3215 Garland Street  
Midland, Michigan 48640(US)  
Inventor: **Morgan, Alice E. R.**  
4610 Bristol Court  
Midland, Michigan 48640(US)

(74) Representative: **Smulders, Theodorus A.H.J.,  
Ir. et al**  
Vereenigde Octrooibureaux Nieuwe Parklaan  
107  
NL-2587 BP 's-Gravenhage(NL)

(54) **Ribosome-inactivating proteins, inactive precursor forms thereof, a process for making and a method of using.**

(57) The present invention is directed to ribosome inactivating proteins. The proteins are characterized by being in a single chain proRIP inactive form that can be converted into an active form by cleavage with proteases.

EP 0 466 222 A1

Ribosome-inactivating proteins (RIPs) are plant proteins that are capable of catalytically inactivating eukaryotic ribosomes and are consequently extremely potent inhibitors of eukaryotic protein synthesis. RIPs have been divided into two classes: type 1 and type 2 RIPs (see Barbieri and Stirpe (1982), Cancer Surveys, 1:489-520). There is significant amino acid sequence homology between members of both type 1 and type 2 RIPs, and with the bacterial Shiga and Shiga-like toxins which also have the same mechanism of action (see Hovde *et al.* (1988), Proc. Natl. Acad. Sci. USA, 85:2568-2572).

Type 2 RIPs consist of two polypeptides: an RIP (or A-chain) which is covalently attached via a disulfide bond to a lectin-like protein (or B-chain) with an affinity for cell surface carbohydrates. The B-chain binds to the cell surface and facilitates subsequent cellular internalization of the RIP A-chain moiety, which results in rapid inactivation of protein synthesis and cell death. Type 2 RIPs are therefore extremely potent cytotoxins and animal poisons, the best studied example of which is ricin.

In contrast type 1 RIPs, characterized to date, consist of a single polypeptide chain equivalent in activity to that of A-chain RIPs but lacking a covalently attached B-chain. Consequently, they are scarcely toxic to intact cells but retain their extreme potency against cell-free protein translation systems. Typical concentrations that inhibit cell-free protein translation by 50 percent ( $IC_{50}$ ) are 0.5 to 10 ng/ml (0.16 to 33 pM). Until the discoveries detailed hereinbelow, reported type 1 RIPs were a remarkably homogeneous class of basic proteins with Mr values of about 30,000. Type 1 RIPs are found in a great variety of dicot and monocot plants and they may be ubiquitous. They are often abundant proteins in seeds, roots or latex for example. Their *in vivo* function is unclear but it has been proposed that they may be antiviral agents (see Stevens *et al.* (1981), Experientia, 37:257-259) or antifungal agents (see Roberts and Seltrennikoff (1986), Bioscience Reports, 6:19-29).

To date, one article has discussed the presence of an inhibitor of animal cell-free protein synthesis in maize, as well as other cereal crops (see Coleman and Roberts (1982), Biochimica et Biophysica Acta, 696:239-244). The preparation of the maize inhibitor was via ammonium sulfate precipitation and phosphocellulose column chromatography. It is generally believed that the inhibitor isolated from maize was pure. The reported molecular weight of the inhibitor was 23 kiloDaltons (kD), with a reported  $IC_{50}$  of 50 to 100 ng/ml in an ascites cell-free protein synthesis assay.

Where the effect of RIPs on ribosomes has been examined, both type 1 and type 2 RIPs possess a unique and highly specific N-glycosidase activity which cleaves the glycosidic bond of adenine 4324 of the rat liver ribosomal 28S RNA (see Endo (1988), In: Immunotoxins, (ed.) Frankel).

Commercial interest in RIPs has primarily focused on their use in construction of therapeutic toxins targeted to specific cells such as tumor cells by attachment of a targeting polypeptide, most frequently a monoclonal antibody (see Immunotoxins (1988), *supra*). This mimics the binding functionality of the B-chain of type 2 RIPs but replaces their non-specific action with a highly selective ligand. Another recent potential use is in HIV therapy (see U.S. Patent 4,869,903 to Lifson *et al.*, (Genelabs Incorporated and The Regents of the University of California)).

However, while a maize-derived protein synthesis inhibitor, like protein synthesis inhibitors from other *Panicoideae* would appear to be useful for construction of cytotoxic conjugates, no artisan to date has reported to have successfully used a *Panicoideae* RIP. This is somewhat surprising in view of the success achieved with RIPs from non-*Panicoideae* plants, including cereals such as barley (see Lambert *et al.* - (1988), In: Immunotoxins, *supra*). In part, the lack of success to date by skilled artisans in successfully utilizing the maize RIP described by Coleman and Roberts may be attributed to the fact that the protein synthesis inhibitor was relatively uncharacterized.

There is interest in expressing recombinant RIP in commonly employed host eukaryotic cells, because of the capacity to provide correct post-translational processing. However, as RIPs effectively inhibit protein synthesis in eukaryotic cells, a predictable problem is that heterologous expression of an RIP will result in host cell death. Thus, eukaryotic cells are generally not used as recombinant host cells. Although eukaryotic cells may be used in certain circumstances, the RIP must be constructed so as to be secreted prior to the cell experiencing toxicity (see EP 0 335 476 to Gelfand *et al.* (Cetus Corp.)). Therefore, prokaryotic host cells are generally used as hosts, notwithstanding disadvantages such as the inability to glycosylate and properly fold heterologously expressed proteins.

It is thus an object of the invention to provide a method of preparing inactive forms of RIPs, in which an inactive RIP may be expressed by eukaryotic host cells and then converted to an active form.

It is yet another object of the invention to provide the DNA sequence of the gene encoding at least one inactive form of RIP, as well as expression vehicles, host cells and cell cultures containing such DNA sequence.

Other objects and advantages of the present invention will become apparent from the teachings presented hereinafter.

It is to these objects to which the present invention is directed.

In a first aspect, the present invention is directed to a homogeneous protein, termed a proRIP, wherein the protein is incapable of substantially inactivating eukaryotic ribosomes, but which can be converted into a protein, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes, said proRIP having a removable, internal peptide linker sequence.

In a second aspect, the present invention is directed to a homogeneous protein derived from *Panicoideae*, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes.

In a third aspect, the present invention is directed to a homogeneous protein derived from *Panicoideae*, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes, wherein the RIP has  $\alpha$  and  $\beta$  fragments.

In a fourth aspect, the present invention is directed to a homogeneous protein derived from maize, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes, said  $\alpha$  fragment having an amino acid sequence effectively homologous to the following sequence:

KRIVPKFTEIFPVEDANYPYSAFIASVRKDVIKHCTDHKGIFQPVLPPEKKVPELW  
FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
AVNDLAKKKK

and said  $\beta$  fragment having an amino acid sequence effectively homologous to the following sequence:

DPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHGVTLTVTQGGKQVQKWDR  
SKAAFEWADHPTAVIPDMQKLGKDKNEAARIVAL  
VKNQTTAAATAASADNDDEA.

In a fifth aspect, the present invention is directed to a method of converting a protein that is incapable of substantially inactivating eukaryotic ribosomes into a protein that is capable of substantially inactivating eukaryotic ribosomes, said method comprising the following steps:

- a) providing a homogeneous protein derived from *Panicoideae*, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes; and
- b) contacting the proRIP with a cleaving agent capable of deleting the linker to form a protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes.

In a sixth aspect, the present invention is directed to a conjugate comprising a targeting vehicle and a protein derived from *Panicoideae*, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes.

In a seventh aspect, the present invention is directed to a conjugate comprising a targeting vehicle and a protein derived from *Panicoideae*, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes, wherein the  $\alpha\beta$  RIP has  $\alpha$  and  $\beta$  fragments.

In an eighth aspect, the present invention is directed to a conjugate comprising a targeting vehicle and a protein derived from maize, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes, said  $\alpha$  fragment having an amino acid sequence effectively homologous to the following sequence:

KRIVPKFTEIFPVEDANYPYSAFIASVRKDVIKHCTDHKGIFQPVLPPEKKVPELW  
 FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
 5 GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
 AVNDLAKKKK

and the  $\beta$  fragment having an amino acid sequence effectively homologous to the following sequence:

10 DPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHGVTTLTVTQGKQVQKWDR I  
 SKAAFEWADHPTAVIPDMQKLGKDKNEAARIVAL  
 15 VKNQTTAAAATAASADNDDDEA.

In a ninth aspect, the present invention is directed to DNA isolate capable of encoding a protein, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a  
 20 protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes.

In a tenth aspect, the present invention is directed to a DNA isolate encoding a protein, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes, wherein the  $\alpha\beta$  RIP has  $\alpha$  and  $\beta$  fragments.

25 In an eleventh aspect, the present invention is directed to a DNA isolate encoding a protein derived from maize, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes, said  $\alpha$  fragment having an amino acid sequence effectively homologous  
 30 to the following sequence:

KRIVPKFTEIFPVEDANYPYSAFIASVRKDVIKHCTDHKGIFQPVLPPEKKVPELW  
 35 FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
 GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
 AVNDLAKKKK

40 and said  $\beta$  fragment having an amino acid sequence effectively homologous to the following sequence:

DPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHGVTTLTVTQGKQVQKWDR I  
 SKAAFEWADHPTAVIPDMQKLGKDKNEAARIVAL  
 45 VKNQTTAAAATAASADNDDDEA.

In a twelfth aspect, the present invention is directed to a DNA isolate encoding a fusion protein capable of substantially inactivating eukaryotic ribosomes, said protein having an amino acid sequence effectively  
 50 homologous to the following sequence:

KRIVPKFTEIFPVEDANYPYSAFIASVRKDVIKHCTDHKGIFQPVLPPEKKVPELW  
 FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
 GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
 5 AVNDLAKKKKDPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHG  
 VTLTVTQGKQVQKWDRISKAAFEWADHPTAVIPDMQKLGKDKN  
 EAARIVALVKNQTTAAAATAASADNDDDEA.

10

In other aspects, the invention is directed to expression vehicles capable of effecting the production of such aforementioned proteins in suitable host cells. It also includes the host cells and cell cultures which result from transformation with these expression vehicles.

A number of aspects of the present invention are further illustrated in the accompanying Drawings, in which:

15

Figure 1 shows a schematic representation of the processing of maize proRIP to the active form.

Figure 2 shows the effect of active maize  $\alpha\beta$  RIP on mammalian cell-free protein synthesis.

The entire teachings of all references cited herein are incorporated by reference.

## 20 Definitions

Nucleic acids, amino acids, peptides, protective groups, active groups and so on, when abbreviated, are abbreviated according to the IUPACIUB (Commission on Biological Nomenclature) or the practice in the fields concerned. The following are examples.

25 The single letter code for amino acids is set forth below:

30

35

Glycine:	G	Phenylalanine:	F
Alanine:	A	Tyrosine:	Y
Valine:	V	Threonine:	T
Leucine:	L	Cysteine:	C
Isoleucine:	I	Methionine:	M
Serine:	S	Glutamic acid:	E
Aspartic acid:	D	Tryptophan:	W
Lysine:	K	Proline:	P
Arginine:	R	Asparagine:	N
Histidine:	H	Glutamine:	Q
Unknown	X		

40 The term "proRIP" means a precursor protein that contains a leader and linker and that is not capable of inactivating eukaryotic ribosomes.

The term "leader" refers to an N-terminal amino acid sequence of a proRIP that need not present in the mature, fully active form of the  $\alpha\beta$  RIP.

The term "linker" refers to an internal amino acid sequence within a proRIP, whereby the linker is of a length and contains residues effective to render the proRIP incapable of catalytically inhibiting translation of a eukaryotic ribosome.

The term "RIP" means a protein that is capable of inactivating eukaryotic ribosomes. The term " $\alpha\beta$  RIP" means a RIP having an  $\alpha$  fragment, which may or may not contain the leader, a  $\beta$  fragment and being capable of substantially inactivating eukaryotic ribosomes.

50 The term "IC<sub>50</sub>" means the concentration of a protein necessary to inhibit protein synthesis by 50 percent in a cell-free protein synthesis assay.

The term "inhibiting amount" refers to the specific ability of RIPs to cause the death or injury of cells against which they are targeted.

The term "target cells" means those cells having ribosomes which the  $\alpha\beta$  RIP of the present invention is capable of inhibiting. The target cells may be present in living organisms or they may be preserved or maintained *in vitro*. The cells may be individual or associated to form an organ. Exemplary target cells include any eukaryotic cell (e.g., mammalian, insect, fungal and plant cells).

The term "targeting vehicle" means a carrier moiety containing a ligand capable of binding to a

receptor of a specific cell or tissue.

"Gene" refers to the entire DNA portion involved in the synthesis of a protein. A gene embodies the structural or coding portion which begins at the 5' end from a translation start codon and extends to a stop codon at the 3' end. It also contains a promoter region, usually located 5' or upstream to the structural coding portion, which initiates and regulates the expression of a structural gene and a 3' nontranslated region downstream from the translated region.

"Expression" refers to a two-part process for the transcription and translation of a gene. The DNA defining the gene is transcribed into a precursor RNA, which is processed to its mature form, messenger RNA (mRNA). During translation, the cell's ribosomes, in conjunction with transfer RNA, translate the RNA "message" into proteins.

#### Preferred Embodiments of the Invention

Surprisingly, it has been discovered that studied members of *Panicoideae* contain  $\alpha\beta$  RIP and proRIP. *Panicoideae* is a subfamily of *Gramineae* (order) and *Graminaceae* (family). The subfamily *Panicoideae* contains three tribes: *Maydeae* (e.g., *Tripsacum*, *Coix*, *Euchlaena* and *Zea*), *Andropogoneae* (e.g., *Sorghum*) and *Paniceae*. For further taxonomic information, see Arber (1934), *The Gramineae, A Study of Cereal, Bamboo and Grass*, Cambridge University Press, p 410-411.

The present invention pertains to proteins which are derived from a plant within the subfamily *Panicoideae*. As taught herein, proteins obtained from various plants within the subfamily *Panicoideae* have shown antigenic cross reactivity (i.e., showing evidence of proRIP in *Panicoideae* as well as  $\alpha$  and  $\beta$  fragments of an  $\alpha\beta$  RIP).

By "derived" from a plant within the subfamily *Panicoideae* means a protein that is effectively homologous, as defined below, with a proRIP or  $\alpha\beta$  RIP from *Panicoideae*, regardless of the manner in which the protein is produced. Given the present teachings it now becomes possible to prepare generally homogeneous proRIP and  $\alpha\beta$  RIP exclusive of irrelevant proteins and contaminants naturally associated therewith in the cellular environment or in extracellular fluids. For example, a substantially pure protein will show constant and reproducible characteristics within standard experimental deviations for parameters such as the following: molecular weight, chromatographic behavior, and such other parameters. The term, however, is not meant to exclude artificial or synthetic mixtures of a protein with other compounds. The term is not meant to exclude the presence of minor impurities which do not interfere with the biological activity of the protein and which may be present, for example, due to incomplete purification.

Both the proRIP and  $\alpha\beta$  RIP may be purified directly from mature and germinating seeds and developing kernels of plants within the subfamily. Generally, the purification of the *Panicoideae*  $\alpha\beta$  RIP and proRIP may be accomplished as follows.

Seeds or immature kernels of plants within the subfamily *Panicoideae* may be homogenized to disrupt the individual seeds or kernels. This can be accomplished by any type of commercially available homogenizer.

The *Panicoideae* proRIP and/or  $\alpha\beta$  RIP may be purified from the homogenization product by any appropriate protein purification technique. Exemplary techniques include gel filtration chromatographic techniques, such as conventional liquid chromatography, ion exchange chromatography, high performance liquid chromatography, and reverse phase chromatography.

Upon purification, the *Panicoideae* proRIP will have insignificant ribosome inactivating ability relative to its corresponding  $\alpha\beta$  RIP. For example, maize proRIP has an  $IC_{50}$  of greater than 10 micrograms per milliliter ( $\mu\text{g/ml}$ ) in a cell-free protein synthesis assay. The maize  $\alpha\beta$  RIP has an  $IC_{50}$  of about 1 nanogram per milliliter ( $\text{ng/ml}$ ) in a mammalian cell-free protein translation assay.

The maize proRIP has a molecular weight of about 34 kD, as determined by SDS-PAGE (see Laemmli (1970), *supra*), and will move as a single peak on ion exchange chromatography. Homogeneous maize  $\alpha\beta$  RIP will comprise two associated fragments, an  $\alpha$  and  $\beta$  fragment, having molecular weights of approximately 16.5 kD and 8.5 kD, respectively (as determined by sodium dodecyl sulfate polyacrylamide-gel electrophoresis (SDS-PAGE) (see Laemmli ((1970), *Nature*, 22:680-685). The homogeneous protein will exhibit two dissociated peaks on reverse phase chromatography, and a single associated peak on ion exchange chromatography. Polyclonal anti-sera against each fragment both crossreact with a polypeptide present in maize kernels having a molecular weight of about 34 kD as determined by SDS-PAGE. This demonstrates that the two fragments of the maize  $\alpha\beta$  RIP are in fact derived from a common precursor (i.e., the maize proRIP).

The maize proRIP amino acid sequence (as set forth in Table 1) contains four sequence subsegments: (1) a leader sequence, from residues 1 to 16, (2) an  $\alpha$  fragment, from residues 17 to 156, (3) an internal



linker sequence, from residues 157 to 182, and (4)  $\alpha$   $\beta$  fragment, from residues 183 to 301.

TABLE 1

5

Nucleotide sequence and deduced amino acid sequence of maize proRIP cDNA.

	1	GAA TTC GGC ACG AGC AAA GAG AAG GGA ATG GGC GAG ATA ACC CTA GAG CCG	Met Ala Glu Ile Thr Leu Glu Pro
10	32	AGT GAT CTT ATG GCG CAA ACA AAC AAA AGA ATA GTG CCA AAG TTC ACT GAA	Ser Asp Leu Met Ala Gln Thr Asn Lys Arg Ile Val Pro Lys Phe Thr Glu
	64	ATC TTC CCC GTG GAG GAC GCG AAC TAC CCT TAC AGC GCG TTC ATC GCG TCG	Ile Phe Pro Val Gln Asp Ala Asn Tyr Pro Tyr Ser Ala Phe Ile Ala Ser
15	128	CTC CCG AAA GAC GTG ATC AAA CAC TCG ACC GAC CAT AAA GCG ATC TTC CAG	Val Arg Lys Asp Val Ile Lys His Cys Thr Asp His Lys Gly Ile Phe Gln
	200	CGC GTG CTG CCA CCG GAG AAG AAG GTC CCG GAG CTA TGG TTC TAC ACA GAG	Pro Val Leu Pro Pro Glu Lys Lys Val Pro Glu Leu Trp Phe Tyr Thr Glu
20	256	CTC AAA ACT AGG ACC AGC TCC ATC ACG CTC GCC ATA CGC ATG GAC AAC CTC	Leu Lys Thr Arg Thr Ser Ser Ile Thr Leu Ala Ile Arg Met Asp Asn Leu
	320	TAC CTC CTC GGC TTC AGG ACC GCG GCG GCG GTC TGG TGG GAG TTC GCG AAG	Tyr Leu Val Gly Phe Arg Thr Pro Gly Gly Val Trp Trp Glu Phe Gly Lys
25	384	GAC GCG GAC ACC GAC CTC CTC GCG GAC AAC CCG AGG TGG CTC GCG TTC GCG	Asp Gly Asp Thr His Leu Leu Gly Asp Asn Pro Arg Trp Leu Gly Phe Gly
	448	GCG AGG TAC GAG GAC CTC ATC GCG AAC AAG GGT CTC GAG ACC CTC ACC ATG	Gly Arg Tyr Gln Asp Leu Ile Gly Asn Lys Gly Leu Glu Thr Val Thr Met
30	512	GCG GCG GCG GAA ATG ACC AGG GCG CTC AAC GAC CTC GCG AAG AAG AAG AAG	Gly Arg Ala Glu Met Thr Arg Ala Val Asn Asp Leu Ala Lys Lys Lys Lys
	576	ATG GCG ACA CTC GAG GAG GAG GAG GTC AAG ATG CAG ATG CAG ATG CCG GAG	Met Ala Thr Leu Glu Glu Glu Glu Val Lys Met Gln Met Gln Met Pro Glu
35	640	GCG GCT GAT CTC GCG GCG GCG GCA GCG GCT GAC CCA CAG GCG GAC AGG AAG	Ala Ala Asp Leu Ala Ala Ala Ala Ala Asp Pro Gln Ala Asp Thr Lys
	704	AGC AAG CTC CTC AAG CTC CTC CTC ATG GTC TCG GAG GCG CTC CCG TTC AAC	Ser Lys Leu Val Lys Leu Val Val Met Val Cys Gln Gly Leu Arg Phe Fun
40	768	ACC CTC TCG CCG ACC CTC GAC GCG GCG TTC AAC AGC CAG GAC GCG CTC ACC	Thr Val Ser Arg Thr Val Asp Ala Gly Phe Asn Ser Gln His Gly Val Thr
	832	TTC ACC CTC ACC GAG GCG AAG CAG CTC CAG AAG TCG GAC AGG ATC TTC AAG	Leu Thr Val Thr Gln Gly Lys Gln Val Gln Lys Trp Asp Arg Ile Ser Lys
45	896	GCG GCG TTC GAG TCG GCT GAC GAC CCG ACC GCT CTC ATC CCG GAC ATC CAG	Ala Ala Phe Gln Trp Ala Asp His Pro Thr Ala Val Ile Pro Asp Met Gln
	960	AAG CTC GCG ATC AAG GAT AAG AAC GAA GCA GCG AGG ATC GTC GCG CTC GTC	Lys Leu Gly Ile Lys Asp Lys Asn Tle Ala Ala Arg Ile Val Ala Lys Val
50	1024	AAG AAT GAA ACT ACT GCG GCT ACC GCT ACT GCT GCG AGT GCT GAC AAC GAC	Lys Asn Gln Thr Thr Ala Ala Ala Ala Thr Ala Ala Ser Ala Asp Asn Asp
	1088	TAC GAC GAG GCG TGA TGA ATG GAA GGA GAC ATC ATG ATC TCG TCG TCG ACT	Asp Asp Glu Ala End
55	1152	GAA TTA CTA TGT TCG TAT ACA TAT AAA TAC ACC CCG GCT ACC GCG TGT TCG	
	1216	CTA TAC GGT CCA TAA TGT ACC GAG TAA ATT TTA AAC TAC TGT TGT GCG	
	1272	AAT TC	

The net charges of these polypeptides are as follows: leader -3;  $\alpha$ , +10; linker, -5;  $\beta$ , +1. Removal of the leader and linker results in a dramatic change in net charge from the maize proRIP (+3) to maize  $\alpha\beta$  RIP (+11). Additionally, the proRIP isolated from maize has an observed pI of 6.5 which agrees well with

the value of 6.1 derived from the deduced amino acid sequence. The pI of the active maize  $\alpha\beta$  RIP is 2.9, compared to the calculated value from the deduced amino acid sequence of 9.06 (i.e., after deletion of the acidic leader, and linker sequences). Experimental data further suggests that the native maize  $\alpha\beta$  RIP has a few amino acids removed from the carboxy-terminal end. Thus, the maize  $\alpha\beta$  RIP has a basic pI, which is consistent with the pI of other RIPs.

A comparison of the maize proRIP with that of barley, a monocot, as set forth in Table 2 below. The upper sequence shows maize  $\alpha\beta$  RIP and the lower sequence barley RIP, as taught by Asano *et al.* (1986), *Carlsberg Res. Commun.*, 51:129. Identical residues are denoted by a solid line, conservative substitutions by a dotted line, and dashes indicate insertions to maximize homology. Residues are numbered on the left.

TABLE 2

## Comparison of maize RIP and barley RIP amino acid sequences.

1	MAEITLPSDLMAQTNKRIVPKFTEIFPVEDANYPYSAFIASVRKDVIKHCTDHKGIFQPV	
20	-----AAKMAKNVDKPLFTATFNVCASSADYATFIAGIRNKLRNPA--HFSHNEPV	
61	LFP-EKKVPEL-WFYTELKTR-TSS-ITLAIRMDNLYLVGFRTPGGVWWEFGKDGDTLLG	
80	LFPVEPNVPPSRWFHVVLKASPTTSAGLTALAIRADNIYLEGFKSSDGTWELTPGLIPGAT-	
25	119 ENPRWLGGGGRYQDLIGNKGLT-VTMGRAEMTRAVNDLAKKKKMATLEEEVKNQNMPE	
110	-----IVGFGGTYRDLLEGDTTKLTNVALGRQQLDAVTALHGRTK-----	
197	AADLAAAAAADPQADTKSKLVKLVVMVCEGLRFNTVSRIVDAGFNSQHGVTLT-----VTQG	
190	-----ADKASGFKQQQAREAVTLLLMVNEATRFQTVSGFVAGLLHPKAVEFKSGKIGNEMK	
236	KQVQKWDRIKAAFEWADHPTAVTPDMQKLGKDKNEAARIVALVKNQTTAAATAASADN	
35	197 AQVNGWQQLS-AALLK-----IDVKPPPCKSPAKFTPIKMGVRTAEQ-----AATLGILLF	
237	DDDEA	
259	WVWFGGLTVAKALELPHASGGK	

As set forth in Table 2, there is an overall homology of 28 percent (34 percent including conservative substitutions) between the maize  $\alpha\beta$  RIP and barley RIP. However, the unique nature of the linker region of maize proRIP is clearly shown by the resulting gap that has been introduced in the published barley sequence to maintain homology. A lower, but significant, degree of homology is seen when the maize proRIP sequence is compared to that of ricin A-chain (as set forth in Table 3 below). The upper sequence is maize  $\alpha\beta$  RIP and the lower sequence is ricin A, as taught by Lamb *et al.* (1985), *Eur. J. Biochem.*, 148:265. Identical residues are denoted by a solid line and conservative substitutions by a dotted line, dashes indicate insertions to maximize homology. Residues are numbered on the left, the numbering of the ricin sequence corresponds to that of the mature protein.

TABLE 3

Comparison of maize RIP and ricin A chain amino acid sequences.

5

10

15

20

25

```

1  -----MAEITLEPSDLMAQTNKRIVPKFTEIFPVEDANYPYSAFIASVRKDVIKHCT
   : : : | : : : : : | | | |
-24 MYAVATWLCFGSTSGWSFTLEDNNIFPKQYPIINFTTAGATVQSYTNFIRAVRGRLLTGAD
10 62 DHKGIFQPVLPPEKKVPELWF-YTELKTR-TSSITLAIRMDNLYLVGFRTPGGVWWEFGK
   : : : | : : : | : : : | : : : | : : : |
38 VR--HEIPVLPNRVGLPINQRFILVELSNHAEISVTIALDVTNAYVVGYRAGNSAYFFHPD
110 DGDTHLLGDNPR-----WLGFGGGRYQDLIGNKGL--ETVTMGRAEMTRAVNDLAKKKKM
15 : : : | : : : | : : : | : : : | : : : |
97 NQEDAEAITHLFTDVQNRYTFAFGGNYDRLEQLAGNLRENIELGNGPLEEAI SALYYYST-
164 ATLEEEEVKMQMPEAADLAAAAAADPQADTKSKLVKLVVMVCEGLRFNTVSRITVDAGFN
   : : : | : : : | : : : | : : : | : : : |
20 157 -----GGTQLPTLARSFIICIQMISEAARFQYIEGEMRTRIR
224 SQHGVTLTVTQGKQVQKWDRIKAAFEWADHPTAVIPDMQKLG I-KDKNEAARIVALVKNO
   : : : | : : : | : : : | : : : | : : : |
194 YNRRSAPDPFSVITLENSWGRSLTAIQESNQGAFA SPIQLQRRNGSKFSVYDVSILIPIAL
25 284 TTAATAASADNDDEA
   : : : | : : : | : : : | : : : | : : : |
255 MVYRCAPPPSQF

```

30 As set forth in Table 3, a gap was again introduced in the published ricin A sequence to maximize homology corresponding to the linker region of the maize proRIP.

Further comparison of the maize proRIP sequence with published full-length sequences of other non-*Panicoideae* RIPs indicate that there are four regions of significant homology between these proteins (as set forth in Tables 4a and 4b below).

35

TABLE 4A

40

Alignment of the N-terminal amino acid sequence of the maize RIP 16.5 kD polypeptide with the N-termini of RIPs from other sources.

45

50

55

```

45 Maize RIP      AQTNKRIVPKFTEIF-PVEDANYPYSAFIASVRKDVIK
   Barley RIP    AAKMAKNVDKPLFTATF-NVQASSADYATFIAGIPNKLRLN
   Ricin A       IFPKQYPIINFTTAGATVQSYTNFIRAVRGRLLT
   Trichosanthin DVSFRLSGATSSSYGVFISNLRKALPN
   Momordin      DVSFRLSGADPRSYGMFIKDLRNALPF
50 Bryodin       DVSFRLSGATTTSYGVFIKNLREALPY
   Gelonin       GLDTVSFSTKGATYITYVNFNLRLVKKLP
   Dodecandrin   VNTIINVGSTTISNYATFMDNLANEAKD
   Pokeweed AP2  N-IVFDYENATPETYSNFLTSLREAVKD
   Saporin 5     VTSITLDLVNPTAGQYSSSFVDKIPNNVKD
55 Saporin 4     VIIYELNLQGTTKAQYSTILKQLRDDIKD
   SLT-1A       KEFTLDFSTAKTYDSLNV-IPSAIGT

```

Table 4A shows the first region and the comparative alignment of the N-terminal amino acid sequence of the maize  $\alpha\beta$  RIP  $\alpha$  fragment with the N-terminal sequences of RIPs from other sources. The sequences are taken from: barley (see Asano *et al.* (1986), *supra*); ricin A-chain (see Lamb *et al.* (1985), *supra*); dodecandrin (see Ready *et al.* (1985), *Biochem. Biophys. Acta*, 791:314); pokeweed anti-viral protein 2 (AP2) (see Bjorn *et al.* (1985), *Biochim. Biophys. Acta*, 790:154); Shiga-like toxin 1A (SLT-1A) (see Calderwood *et al.* (1987), *Proc. Nat. Acad. Sci. USA*, 84:4364); and  $\alpha$ -trichosanthin, momordins, bryodin, gelonin, dodeoandrin, pokeweed antiviral protein-2, saporin 5, and saporin 4 (see Montecucchi *et al.* (1989), *Int. J. Peptide Res.*, 33:263). Positions showing homology in four or more sequences are noted by solid lines (showing identical residues) or dotted lines (showing conservatively substituted residues).

TABLE 4B

**Alignment of maize RIP with regions of homology in the amino acid sequences of other RIPs.**

20	Maize RIP	84	<u>ITLAIRMONLYLVGE</u>	201	<u>LVVMVCEGLRENTYS</u>	237	<u>QVOK-WDRISKA</u>
	Barley RIP	77	<u>LTALIRADNIXLEGE</u>	168	<u>LLLMVNEATREFOTYS</u>	208	<u>QVNG-WODLSAA</u>
	Ricin A-chain	70	<u>VTALADVTNAXVVGX</u>	171	<u>CIOMISEAARFOYLE</u>	207	<u>LENS-HGRISTA</u>
	Saporin-6	62	<u>YSLGLKRONLYVVGX</u>	170	<u>AIQMTAEAAERFYIQ</u>	205	<u>EVN--WKKISTA</u>
	Abrin A-chain	64	<u>IEVGIDVTNAXVVGX</u>	159	<u>LIQVASEAARERYIS</u>	194	<u>LENN-WDNLRGV</u>
25	SLT-1A	62	<u>MIDSGSGDNLEAVDV</u>	183	<u>FYTVTAEALRFOIQR</u>	222	<u>TLN--WGRLLSV</u>
	Trichosanthin	58	<u>DEVALDVTNV...</u>	161	<u>LIQSTSEAAARYKFIE</u>	196	<u>LENSLWLALSKQ</u>
		77	<u>YLMGY</u>				

Table 4b shows that the other three regions are internally oriented. Table 4b specifically shows the alignment of maize  $\alpha\beta$  RIP with regions of homology in the amino acid sequences of other RIPs. The sequences are available from the following references: barley (see Asano *et al.* (1986), *supra* and Leah *et al.* (1991), *J. Biol. Chem.*, 266:1564-1573); ricin A-chain (see Lamb *et al.* (1985), *supra*); abrin A-chain (see Funatsu *et al.* (1988), *Agric. Biol. Chem.* 52:1095); saporin-6 (see Benatti *et al.* (1989) *Eur. J. Biochem.*, 183:465); Shiga-like toxin 1A (SLT-1A) (see Calderwood *et al.* (1987), *supra*); and  $\alpha$ -trichosanthin (see Xuejun and Jiahuai (1986), *Nature*, 321:477; Chow *et al.* (1990), *J. Biol. Chem.*, 265:8670-8674 and Maragonore *et al.* (1987), *J. Biol. Chem.*, 262:11628-11633). Positions showing identity or conservative substitutions in four or more sequences are underlined, dashes indicate insertions to maximize homology. Vertical lines indicate residues that are conserved in all seven sequences. The starting amino acid of each sequence is indicated (note that trichosanthin contains an insertion sequence at residues 67 to 76).

The sequences and partial sequences of various additional Type I RIPs are set forth in the following articles: luffin-A (see Islam *et al.* (1990), *Agric. Biol. Chem.*, 54:2967-2978; mirabilis antiviral protein (see Habuka *et al.* (1989) *J. Bio. Chem.*, 264:6629-6637; trichokirin, (see Casellas *et al.* (1988), *Eur. J. Biochem.*, 176:581-588; momordins (see Barbieri *et al.* (1980), *Biochem. J.*, 186:443-452; dianthins (see Reisbig and Bruland (1983), *Arch. Biochem. Biophys.*, 224:700-706; saporins (see Maras *et al.*, (1990), *Biochem. Intl.*, 21:831-838 and Lappi *et al.* (1985), *Biochem. Biophys. Res Commun.*, 129:934-942; and momorcochin-S (see Bolognesi *et al.* (1989), *Biochim. Biophys. Acta*, 993:287-292).

Other Type I and Type II RIPs have also been purified to homogeneity and these include; momorcharins (see Yeung *et al.* (1986), *Int. J. Peptide Res.*, 28:518-524; tritins (see Roberts and Stewart (1979), *Biochem.*, 18:2615-2621); rye (see Coleman and Roberts (1982), *Biochim. Biophys. Acta*, 696:239-244; agrostins and *Hura crepitans* (see Stirpe *et al.* (1983), *Biochem. J.*, 216:617-625; *Asparagus officinalis* - (see Stirpe *et al.* (1983), *Biochem. J.*, 216:617-625; *Cucumis melo* (see Ferreras *et al.* (1989), *Biochem Intl.*, 19:201-207; *Cucurbitaceae* (see Ng *et al.*, *Int. J. Biochem.*, 21:1353-1358; *Petrocoptis* (see Ferreras *et al.*, *Cell. Molec. Biol.*, 35:89-95); volkensin-a (see Barbieri *et al.* (1984), *FEBS Lett.*, 171:277-279; viscumin-a (see Olsnes *et al.* (1982), *J. Biol. Chem.*, 257:13263-13270; modeccin-a (see Gasperi-Campani (1978), *Biochem. J.*, 174:491-496); *Momordia charantia* lectin-a (see Lin *et al.* (1978), *Linn. Toxicon.*, 16:653-660); *Phoradendron californicum* lectin-a (see Franz *et al.* (1989), *FEBS Lett.*, 248:115-118).

Proteins from the following other plants have also been shown to possess ribosome inactivating activity:

*Stellarea holostea*, *Lychnis flos-cuculi*, *Hordeum murinum*, *Aegylops geniculata*, *Euphorbia serrata*, *Capsella bursapastoris*, *Muscari comosum* (see Merino *et al.* (1990), *J. Exp. Botany*, 41:67-70); and proteins from *Croton tiglium* and *Jatropha curcas* (see Stirpe *et al.* (1976), *Biochem. J.*, 156:1-6).

When the internal linker sequence of the proRIP is removed, the  $\alpha\beta$  RIP has significant ribosome inactivating activity. The activity has been found to be significant regardless of whether the leader sequence has been removed (e.g., by recombinant methods). However, the proRIP is most active when the leader sequence is also removed and probably when C-terminus residues are also removed. In nature, it is thought that the linker is cleaved by endogenous proteases released by germinating seeds. Significantly, it has been discovered that the linker may also be cleaved *in vitro* by certain proteases, e.g., papain, to yield active maize  $\alpha\beta$  RIP from the precursor. While not intended to be bound by theory, it is thought that papain likely mimics the effect of endogenous endoproteases released during germination.

It appears that, after removal of the internal linker, the two fragments of the processed polypeptide are held together by noncovalent forces. That is, the association of the two polypeptide chains does not depend upon interchain disulfide bonds or the formation of a peptide bond between the fragments.

Although not intended to be bound by theory, it is believed that the linker forms an external loop with exposed amino acid residues that are susceptible to proteolysis. Support for this suggestion comes from the alignment of the amino acid sequence of the maize proRIP with that of ricin A chain, the three dimensional structure of which is known (see Montfort *et al.* (1987), *J. Biol. Chem.*, 262:5398). The Glu 177, Arg 180, Asn 209 and Trp 211 of ricin A have been implicated in the active site region of the molecule (see Robertus (1988), *In: Immunotoxins, supra*).

Based on this alignment, homologous residues of maize  $\alpha\beta$  RIP can be positioned within the three dimensional structure of ricin A chain. The superimposed structures indicate that the C-terminal lysine of the  $\alpha$  fragment (at residue 162) is in corresponding alignment with an externally positioned threonine (at residue 156) of the ricin A-chain. Also, the N-terminal alanine of the  $\beta$  fragment (at residue 189) is in corresponding alignment with an externally positioned glycine (at residue 157) of the ricin A-chain.

The present invention is intended to include the construction of  $\alpha\beta$  RIP and proRIP forms of any RIP. As set forth in Tables 4a and 4b, RIPs for which a full-length sequence has been determined contain regions with significant homology. Similarly the N-terminal sequence similarities in an even greater number of RIPs have been compared (set forth in Tables 4a and 4b). It is likely that these regions have particular effect upon the function of the respective RIP.

An RIP having known amino acid sequences may now be altered into inactive, proRIP forms by the insertion of a linker. Based on the information deduced from the maize system set forth herein, it now becomes possible to engineer inactive forms of any RIP having a three dimensional structure similar to the three dimensional structure of ricin A chain. Cleavage of the linker will result in an  $\alpha\beta$  RIP not heretofore found in nature.

The art has discussed the methodology for modifying the three dimensional structure of proteins (see, for example, Van Brunt (1986), *Biotechnology*, 4:277-283). The first step involves selecting plausible sites on the RIP between which the linker may be inserted. One of those sites is the exposed amino acid residues surrounding residue 156 of ricin A-chain or its equivalent in other RIP sequences. Thus, the present invention is intended to encompass the insertion of a peptide linker in those sequences, provided that the insertion of the linker substantially reduces the ribosome inactivating ability of the RIP. By "substantially reduce" is meant that the insertion of a cleavable linker into an active RIP lowers the IC<sub>50</sub> value of the resultant protein by at least 10 fold, preferably 100-fold, and more preferably 1000-fold.

As stated previously, ricin A-chain has been shown to have sequence homology to many single chain RIPs. The RIPs set forth in Tables 4a and 4b are intended for exemplification purposes only. RIPs characterized in the future that meet the above criteria are also considered to be a part of this invention.

Generally, the linker may be of a length, may be of an amino acid sequence, and may be internally positioned so as to substantially reduce the ribosome inactivating activity of the RIP.

Obviously, since the *Panicoideae* linker(s) is the only known RIP linker found in nature, it is expected that such an amino acid sequence will logically be a primary candidate for insertion into other RIPs. However the present invention is intended to encompass linkers having effectively homologous sequences to a selected maize linker. The factors to be considered in synthetically preparing effectively homologous linkers for  $\alpha\beta$  RIPs generally are the same as set forth above for selecting effectively homologous linkers for a selected maize linker.

Any of a variety of procedures may be used to clone proRIP-encoding gene sequence. One method for cloning the proRIP gene sequence entails determining the amino acid sequence of the proRIP molecule. To accomplish this task proRIP or  $\alpha\beta$  RIP protein may be purified (as described above), and analyzed to determine the amino acid sequence of the proRIP or  $\alpha\beta$  RIP. Any method capable of elucidating such a

sequence can be employed, however, Edman degradation is preferred. The use of automated sequencers is especially preferred.

It is possible to synthesize *in vitro* the proRIP and  $\alpha\beta$  RIP from their constituent amino acids. A suitable technique includes the solid phase method (see Merrifield (1963), *J. Amer. Chem. Soc.*, 85:2149-2154; and Solid Phase Peptide Synthesis (1969), (eds.) Stewart and Young). Automated synthesizers are also available.

The peptides thus prepared may be isolated and purified by procedures well known in the art (see *Current Protocols in Molecular Biology* (1989), (eds.) Ausubel, *et al.*, Vol. 1) and Sambrook *et al.* (1989), *Molecular Cloning: A Laboratory Manual*).

Using the amino acid sequence information, the DNA sequences capable of encoding them are examined in order to clone the gene encoding the proRIP. Because the genetic code is degenerate, more than one codon may be used to encode a particular amino acid.

Although it is possible to determine the entire amino acid sequence of the proRIP or  $\alpha\beta$  RIP, it is preferable to determine the sequence of peptide fragments of the molecule, and to use such sequence data to prepare oligonucleotide probes which can be used to isolate the entire proRIP gene sequence. The proRIP peptide fragments can be obtained by incubating the intact molecule with cyanogen bromide, or with proteases such as papain, chymotrypsin or trypsin.

Using the genetic code one or more different oligonucleotides can be identified. The probability that a particular oligonucleotide will, in fact, constitute the actual proRIP encoding sequence can be estimated by considering abnormal base pairing relationships and the frequency with which a particular codon is actually used (to encode a particular amino acid) in eukaryotic cells. Using these rules, a single oligonucleotide, or a set of oligonucleotides, that contains a theoretical "most probable" nucleotide sequence capable of encoding the proRIP or  $\alpha\beta$  RIP peptide sequences may be identified.

The oligonucleotide, or set of oligonucleotides, containing the theoretical "most probable" sequence capable of encoding the proRIP gene fragments may be used to identify the sequence of a complementary oligonucleotide, or set of oligonucleotides, which is capable of hybridizing to the "most probable" sequence, or set of sequences. An oligonucleotide containing such a complementary sequence can be employed as a probe to identify and isolate the toxin gene (see Sambrook *et al.* (1989), *supra*).

By hybridizing an oligonucleotide having a sequence complementary to the "most probable" gene sequence, one obtains a DNA molecule (or set of DNA molecules), capable of functioning as a probe to identify and isolate the proRIP gene.

The present invention also relates to DNA sequences that encode recombinant proRIP and  $\alpha\beta$  RIP. The recombinantly-produced proRIP and  $\alpha\beta$  RIP share the following properties with the proRIP and  $\alpha\beta$  RIP isolated from nature and characterized according to the teachings herein: (1) portions of the amino acid sequence deduced from the nucleotide sequence are equivalent to amino acid sequences obtained directly from nature; (2) the polypeptide is recognized by anti-RIP antibodies; (3) the molecular weight of the proRIP and  $\alpha\beta$  RIP polypeptides encoded corresponds with the naturally occurring proteins; (4) each proRIP protein is convertible to an  $\alpha\beta$  RIP; and (5) each proRIP and  $\alpha\beta$  RIP protein exhibits relatively equivalent ribosome inactivating activity.

The process for genetically engineering the proRIP or  $\alpha\beta$  RIP according to the invention is facilitated through the cloning of genetic sequences which are capable of encoding the proRIP or  $\alpha\beta$  RIP, or effectively homologous variants thereof as discussed below, and through the expression of such genetic sequences. As used herein, the term "genetic sequences" is intended to refer to a nucleic acid molecule (preferably DNA). Genetic sequences which are capable of encoding the toxin may be derived from a variety of sources. These sources include genomic DNA, cDNA, synthetic DNA, and combinations thereof.

Cells containing the desired sequence may be isolated, and genomic DNA fragmented by one or more restriction enzymes. The genomic DNA may or may not include naturally-occurring introns. The genomic DNA digested with selected restriction endonucleases yields fragments containing varying numbers of base pairs (bp).

Specifically comprehended as part of this invention are genomic DNA sequences encoding allelic variant forms of the proRIP gene which may include naturally occurring introns. The allelic gene may be derived using a hybridization probe made from the DNA or RNA of the proRIP gene as well as its flanking regions. "Flanking regions" are meant to include those DNA sequences 5' and 3' of the proRIP coding sequences.

The DNA isolate encoding the proRIP gene may also be obtained from a cDNA library. The mRNA may be isolated from a suitable source employing standard techniques of RNA isolation, and the use of oligo-dT cellulose chromatography to enrich for poly-A mRNA. A cDNA library is then prepared from the mixture of mRNA using a suitable primer, preferably a nucleic acid sequence which is characteristic of the desired

cDNA. A single stranded DNA copy of the mRNA is produced using the enzyme reverse transcriptase. From the single stranded cDNA copy of the mRNA, a double-stranded cDNA molecule may be synthesized using either reverse transcriptase or DNA polymerase.

It is also possible to use primers to amplify the DNA from cells of appropriately prepared seeds and immature kernels by the polymerase chain reaction (PCR). PCR involves exponentially amplifying DNA *in vitro* using sequence specified oligonucleotides. (see Mullis *et al.* (1987), *Math. Enz.*, 155:335-350; Horton *et al.* (1989), *Gene*, 77:61; and PCR Technology: Principles and Applications for DNA Amplification, (ed.) Erlich (1989).

The DNA encoding the proRIP or  $\alpha\beta$  RIP may be chemically synthesized by manual procedures, e.g., the phosphotriester and phosphodiester methods (see Caruthers (1983), In: *Methodology of DNA and RNA*, (ed.) Weissman); or automated procedures e.g., using diethylphosphoramidites are used as starting materials (see Beaucage *et al.* (1981), *Tetrahedron Letters*, 22:1859-1962). The DNA may be constructed by standard techniques of annealing and ligating fragments or by other methods.

Thereafter, the desired sequences may be isolated and purified by procedures well known in the art (see Current Protocols in Molecular Biology (1989), *supra*) and Sambrook *et al.* (1989), Molecular Cloning: A Laboratory Manual.

The nucleotide sequence of the maize proRIP cDNA and the deduced amino acid sequence of such the corresponding maize proRIP is set forth in Table 1.

One need not be confined to the amino acid sequences of proRIP and  $\alpha\beta$  RIP found in nature. Thus, it is possible to selectively produce both proRIP and  $\alpha\beta$  RIP via the application of recombinant DNA technology. This in turn allows the production of sufficient quality and quantity of material to create novel forms of the protein unimpeded by the restriction necessarily inherent in the isolation methods involving production and extraction of the protein from natural sources.

Recombinant procedures make possible the production of effectively homologous proteins possessing part or all of the primary structural conformation and/or one or more of the biological properties of the  $\alpha\beta$  RIP. For purposes of this invention, an amino acid sequence is effectively homologous to a second amino acid sequence if at least 70 percent, preferably at least 80 percent, and most preferably at least 90 percent of the active portions of the amino acid sequence are identical and retains its intended function. Thus more importantly and critical to the definition, an effectively homologous sequence to the  $\alpha\beta$  RIP retains the capacity to interact with and inactivate eukaryotic ribosomes. The effectively homologous sequence to the proRIP must retain the capacity to be converted into an  $\alpha\beta$  RIP. That is, the effectively homologous proRIP must have a linker sequence which, when cleaved, will yield a biologically functional  $\alpha\beta$  RIP.

General categories of potentially-equivalent amino acids are set forth below, wherein, amino acids within a group may be substituted for other amino acids in that group: (1) glutamic acid and aspartic acid; (2) lysine, arginine and histidine; (3) hydrophobic amino acids such as alanine, valine, leucine and isoleucine; (4) asparagine and glutamine; (5) threonine and serine; (6) phenylalanine, tyrosine and tryptophan; and (7) glycine and alanine..

It is envisioned that, compared with changes to the  $\alpha$  and  $\beta$  fragments, more significant changes may be made to the proRIP in the leader and linker regions. That is, since the leader and linker sequences are to be cleaved, the length and amino acid residues in their sequences may better be tolerated and considered insignificant, because it will not alter the functionality of the final product.

Further, the linker sequence of the proRIP need not be limited to the sequence set forth in Table 1, above. For example, the length of the linker may be modified, provided that (1) the linker is cleavable, and (2) upon cleavage of the linker the resultant protein has an  $IC_{50}$  value that is at least about 10 times lower than the  $IC_{50}$  value of the protein containing the linker.

Primary criteria for selecting an effectively homologous linker include altering the net charge of the  $\alpha\beta$  RIP (e.g., more acidic); creating a conformational shift in the protein or providing steric hindrance to the active site of the protein.

As noted previously, the maize  $\alpha\beta$  RIP, like other RIPs, is basic. However, the maize proRIP has a slightly acidic pI. Thus, it is preferred that any effectively homologous linker selected for the maize proRIP will be acidic.

The linker should be of a length which, while capable of altering the three-dimensional structure of the protein, when cleaved will permit the protein to retain most of the three dimensional features of the active  $\alpha\beta$  RIP molecule.

To ensure that the linker is cleavable it is generally required that the conformation of the proRIP be such that the linker cleavage sites are readily accessible to a selected cleavage agent.

It is also envisioned that at least one restriction enzyme site may be engineered into the genetic sequence encoding an RIP, allowing DNA sequences encoding various polypeptide linkers to be inserted

into the gene and tested for their ability to create an inactive, yet activatable RIP.

Most commonly, cleavage will be effected outside of the replicative environment, for example, following harvest of microbial culture. Thus, when genetically modifying the proRIP, it may be preferable, in some instances, that the internal linker domain of the proRIP be retained, or altered so as to mimic the manner in which a natural, inactive proRIP liberates its active  $\alpha$  and  $\beta$  fragments.

Any chemical or enzymatic method which recognizes a specific sequence or structure and causes an appropriate cleavage at a selected site may be utilized for the present invention. For example, it may be desirable to design carboxy termini and amino termini of the linker sequences that are subject to cleavage with selected agents. Exemplary such sequences Pro-Xxx-Gly-Pro (where Xxx is unspecified) which is selectively cleaved by collagenase; Ile-Glu-Gly-Arg which is selectively cleaved by Factor Xa; and Gly-Pro-Arg which is selectively cleaved by thrombin (see Nilsson *et al.* (1988), In: Advances in Gene Technology; Protein Engineering and Production, (ed.) Brew *et al.*).

A chemical or enzymatic method may not be suitable if its cleavage site occurs within the active amino acid sequences of the  $\alpha$  and  $\beta$  fragments. That is, cleavage within the native amino acid sequence of the  $\alpha$  and  $\beta$  fragments will generally have a greater likelihood of deleteriously affecting the enzymatic activity of the  $\alpha\beta$  RIP. It is possible to select a specific cleavage sequence of only one amino acid residue so long as that residue does not occur in the amino acid sequences of the  $\alpha$  and  $\beta$  fragments. It is preferred, however, to utilize a specific cleavage sequence which contains two or more amino acid residues, i.e., an extended specific cleavage sequence. This type of sequence takes advantage of the extended active sites of various enzymes. Additionally, by utilizing an extended specific cleavage sequence, it is highly probable that cleavage will only occur at the desired site and not at other sites within the protein.

The cleavage techniques discussed here are by way of example and are but representative of the many variants which will occur to the skilled artisan in light of the specification.

In some instances it may prove desirable to effect cleavage of the proRIP within the cell. For example, an expression vehicle with appropriate promoters may be provided with a DNA sequence coding for enzymes which convert the proRIP to the active form, operating in tandem with the other DNA sequence coding expression of the proRIP.

Further, as is well known, protein sequences may be modified by post-translation processing such as becoming associated with other molecules, for example, glycosides, lipids, or such inorganic ions as phosphate. The ionization status will also vary depending on the pH of the medium or the pH at which crystallization or precipitation of the isolated form occurs. Further, the presence of air may cause oxidation of labile groups, such as -SH. Thus, included within the definition of the proRIP and  $\alpha\beta$  RIP, and fragments, thereof are all such modifications of a particular primary structure, e.g., both glycosylated and non-glycosylated forms, neutral forms, acidic and basic salts, lipid or other associated peptide forms, side chain alterations due to oxidation or derivatization, and any other such modifications of an amino acid sequence which would be encoded by the same genetic codon sequence.

Exemplary techniques for nucleotide replacement include the addition, deletion or substitution of various nucleotides, provided that the proper reading frame is maintained. Exemplary techniques include using polynucleotide-mediated, site-directed mutagenesis, i.e., using a single strand as a template for extension of the oligonucleotide to produce a strand containing the mutation (see Zoller *et al.* (1982), Nuc. Acids Res., 10:6487-6500; Norris *et al.* (1983), Nuc. Acids Res., 11:5103-5112; Zoller *et al.* (1984), DNA, 3:479-488; and Kramer *et al.* (1982), Nuc. Acids Res., 10:6475-6485) and by using PCR, i.e., exponentially amplifying DNA *in vitro* using sequence specified oligonucleotides to incorporate selected changes (see PCR Technology: Principles and Applications for DNA Amplification, Erlich, (ed.) (1989); and Horton *et al.*, *supra*).

By appropriate choice of restriction sites, the desired DNA fragment may be positioned in a biologically functional vector which may contain appropriate control sequences not present in the selected DNA fragment. By "biologically functional" is meant that the vector provides for replication and/or expression in an appropriate host, either by maintenance as an extrachromosomal element or by integration into the host genome. A large number of vectors are available or can be readily prepared, and are well known to skilled artisans.

In general, vectors containing the appropriate promoters, which can be used by the host organism for expression of its own protein, also contain control sequences, ribosome binding sites, and transcription termination sites. Generally, the replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts.

Finally, the vectors should desirably have a marker gene that is capable of providing a phenotypical property which allows for identification of host cells containing the vector.

When expressing the  $\alpha\beta$  RIP, the DNA fragments encoding the  $\alpha$  fragment and the  $\beta$  fragment may be



inserted into separate vectors, or into the same vector. Specifically, when the  $\alpha$  and  $\beta$  fragments are contained in separate vectors, the host cells may be transformed via either cotransformation or targeted transformation techniques.

Construction of suitable vectors containing the desired coding and control sequences may be produced as follows.

Restriction endonucleases may be used as a means for inserting the DNA fragments containing the proRIP gene or  $\alpha\beta$  RIP genes into an appropriate expression vehicle. Exemplary restriction enzymes include *Aat* II, *Bam* HI, *Eco* RI, *Hind* III, *Nde* I, *Spe* I, *Xba* I, *Sac* I, *Bgl* II, *Pst* I, *Sal* I and *Pvu* II.

Cleavage is performed by treating the expression vehicle with a restriction enzyme(s). In general, 10  $\mu$ g vector or DNA fragments is used with 10 units of enzyme in 100  $\mu$ l of buffer solution. Endonuclease digestion will normally be carried out at temperatures ranging from 37 degrees Centigrade (37°C) to 65°C, at a pH of 7 to 9. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturers. Time for the reaction will be from 1 to 18 hours.

After the restriction enzyme digestion is complete, protein may be removed by standard techniques (e.g., extraction with phenol and chloroform). The nucleic acid may be then recovered from the aqueous fraction by standard techniques.

The desired fragment is then purified from the digest. Suitable purification techniques include gel electrophoresis or sucrose gradient centrifugation. The vector and foreign DNA fragments may then be ligated with DNA ligase.

An appropriately buffered medium containing the DNA fragments, DNA ligase, and appropriate cofactors is employed. The temperature employed will be between 4°C to 25°C. When DNA segments hydrogen bond, the DNA ligase will be able to introduce a covalent bond between the two segments. The time employed for the annealing will vary with the temperature employed, the nature of the salt solution, as well as the nature of the sticky ends or cohesive termini. Generally, the time for ligation may be from 5 to 18 hours (see Sambrook *et al.* (1989), *supra*).

Thereafter, the vector constructions may be used to transform an appropriate host cell. Suitable host cells include cells derived from unicellular as well as multicellular organisms which are capable of being grown in cultures or by fermentation.

Various unicellular microorganisms can be used for both cloning and expression. Prokaryotes include members of the Enterobacteriaceae, such as strains of *Escherichia coli*, and *Salmonella*; Bacillaceae, such as *Bacillus subtilis*; *Pneumococcus*, *Streptococcus*, and *Haemophilus influenzae*.

In addition to prokaryotes, eukaryotic cells may be employed. As previously stated, eukaryotic cells have not heretofore been used as recombinant host cells for RIPs. By providing inactive forms of RIPs, the present invention provides skilled artisans with the flexibility to use eukaryotic cells as recombinant hosts. By transforming eukaryotic cells with the proRIP gene, the protein may be expressed at high levels without being toxic to the host cell. Since the protein is lacking in bioactivity pending extra-cellular cleavage, the effect is to enhance the biosafety of the procedure. The proRIP may then be selectively chemically or enzymatically converted to the desired  $\alpha\beta$  RIP.

Exemplary eukaryotic microbes include yeast. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among eukaryotic microorganisms, although a number of other hosts are commonly available.

In addition to eukaryotic microbes, cultures of cells derived from multicellular organisms may also be used as hosts. Examples of useful host mammalian cell lines are Sp2/O, VERO and HeLa cells, Chinese hamster ovary (CHO) cell lines, and W138, BHK, COS-7 and MDCK cell lines.

Other suitable hosts and expression systems are the baculovirus systems maintained in cultured insect cells, e.g., from *Spodoptera frugiperda*.

Finally, cells from and portions of higher plants have been found useful as recombinant hosts, and appropriate control sequences are available for expression in these systems. Suitable plant cells include cells derived from, or seedlings of, tobacco, petunia, tomato, potato, rice, maize and the like.

The expression vehicle may be inserted into the host cell by any suitable method. Conventional technologies for introducing biological material into living cells include electroporation (see Shigekawa and Dower (1988), *Biotechniques*, 6:742; Miller, *et al.* (1988), *Proc. Natl. Acad. Sci. USA*, 85:856-860; and Powell, *et al.* (1988), *Appl. Environ. Microbiol.*, 54:655-660); direct DNA uptake mechanisms (see Mandel and Higa (1972), *J. Mol. Biol.*, 53:159-162; Dityatkin, *et al.* (1972), *Biochimica et Biophysica Acta*, 281:319-323; Wigler, *et al.* (1979), *Cell*, 16:77; and Uchimiya, *et al.* (1982), in: *Proc. 5th Intl. Cong. Plant Tissue and Cell Culture*, A. Fujiwara (ed.), Jap. Assoc. for Plant Tissue Culture, Tokyo, pp. 507-508); fusion mechanisms (see Uchidaz, *et al.* (1980), in: *Introduction of Macromolecules Into Viable Mammalian Cells*, Baserga *et al.* (eds.) *Wistar Symposium Series*, 1:169-185); infectious agents (see Fraley, *et al.* (1986), *CRC Crit.*

Rev. Plant Sci., 4:1-46; and Anderson (1984), Science, 226:401-409; microinjection mechanisms (see Crossway, *et al.* (1986), Mol. Gen. Genet., 202:179-185); and high velocity projectile mechanisms (see EPO 0 405 696 to Miller, Schuchardt, Skokut and Gould, (The Dow Chemical Company). The appropriate procedure may be chosen in accordance with the plant species used.

5 Generally after transformation, the host cells may be grown for about 48 hours to allow for expression of marker genes. The cells are then placed in selective medium and/or screenable media, where untransformed cells are distinguished from transformed cells, either by death or a biochemical property.

The transformed cells are grown under conditions appropriate to the production of the desired protein, and assayed for expression thereof. Exemplary assay techniques include enzyme-linked immunosorbent  
10 assay, radioimmunoassay, or fluorescence-activated cell sorter analysis, immunohistochemistry and the like.

Selected positive cultures are subcloned in order to isolate pure transformed colonies. A suitable technique for obtaining subclones is via the limiting dilution method.

### Uses

15 Essentially all of the uses that the prior art has envisioned for RIPs are intended for the novel  $\alpha\beta$  RIP and proRIP set forth herein (see Immunotoxins (1988), Frankel (ed.); and U.S. Patent 4,869,903 to Lifson *et al.* (Genelabs Incorporated and the Regents of the University of California)).

By providing inactive precursor forms of the  $\alpha\beta$  RIP, it is now possible to provide protein synthesis  
20 inhibitors with uses in practical and improved ways not before possible. The inactive form of the  $\alpha\beta$  RIP offers the additional advantage, over active RIPs, of not being active until removal of the linker sequence. Although the RIP is not toxic to the majority of mammalian cells it is known that RIP may be made specifically cytotoxic by attachment to a targeting vehicle which is capable of binding to and into target cells.

25 Exemplary targeting vehicles include any peptide hormone, growth factor, or other polypeptide cell recognition protein for which a specific receptor exists. A few examples include: antibodies and antibody fragments, lectins, insulin, glucagon, endorphins, growth hormone, melanocyte-stimulating hormone, transferrin, bombesin, low density lipoprotein, luteinizing hormone and asialoglycoprotein that bind selectively to target cells (see Immunotoxins (1988), *supra*). It is well established that conjugates which contain RIP  
30 exhibit maximal cytotoxicity only when the RIP moiety is released from the targeting vehicle.

Since the  $\alpha\beta$  RIP and proRIP does not contain a reactive sulfhydryl group, it may be necessary to modify the proteins using chemical crosslinking reagents in order to link such proRIP and  $\alpha\beta$  RIP to targeting vehicles.

Conjugates of a monoclonal antibody and the  $\alpha\beta$  RIP and proRIP may be made using a variety of  
35 bifunctional protein coupling agents. General examples of such reagents are N-succinimidyl-3-(2-pyridyldithio)propionate, 2-iminothiolane, bifunctional derivatives of imidoesters such as dimethyl adipimidate, active esters such as disuccinimidyl suberate, aldehydes such as glutaraldehyde, bis-azido compounds such as bis(p-diazoniumbenzoyl)ethylenediamine, diisocyanates such as toluene 2,6-diisocyanate, and bis-active fluorine compounds such as 1,5-difluoro-2,4-dinitrobenzene (see, for example, WO 86/05098).

40 In addition, recombinant DNA methodologies may be employed to construct a cytotoxic fusion protein. For a general discussion of chimeric toxins (see, for example, Pastan and FitzGerald (1989), The Journal of Biological Chemistry, 264:15157-15160; and U.S. Patent 4,892,827 to Pastan *et al.* (The United States of America as represented by the Department of Health and Human Services)). These references teach modified *Pseudomonas* exotoxins which comprise a deletion in a receptor binding domain to form a fusion  
45 protein capable of rendering the modified toxin less toxic to selected cells. These references teach DNA sequences encoding the human alpha transforming growth factor ( $\alpha$ -TGF) or human interleukin fused to toxin genes.

### Examples

50 The present invention is illustrated in further detail by the following examples. The examples are for the purposes of illustration only, and are not to be construed as limiting the scope of the present invention. All temperatures not otherwise indicated are Centigrade. All parts and percentages are by weight unless otherwise specifically noted.

#### Example 1: Isolation of Maize $\alpha\beta$ RIP

All steps were performed at 4°C, except for high performance liquid chromatography (HPLC) which was

performed at room temperature. Five hundred grams (500 g) of finely ground mature maize kernels were extracted for at least 2 hour and up to 24 hours (hr) with 1500 ml, 25 mM sodium phosphate, pH 7.2 (PB) + 50 mM sodium chloride. After the extract was strained through several layers of cheesecloth, the protein precipitating between 55 percent and 85 percent ammonium sulfate was collected and redissolved in PB, then dialyzed overnight against the same buffer. The solution was clarified by centrifugation and applied to a 2.5 x 10 centimeter (cm) DE-52 cellulose column equilibrated with PB. The protein passing straight through the column was collected and applied to a Mono S 10/10 column (Pharmacia LKB Biotechnology, Piscataway, NJ) equilibrated with PS, and eluted with a linear gradient of 0 to 200 mM sodium chloride in PB over 90 min at 2 milliliter/ minute (ml/min). Alternatively, the protein can be precipitated with 85 percent ammonium sulfate and dialyzed overnight before applying to the Mono S 10/10 column.

Fractions containing ribosome inactivating protein activity (as measured by rabbit reticulocyte protein synthesis assay, described above) were pooled and concentrated to 0.5 ml in Centricon-10 devices (Amicon, Danvers, MA), and applied to a Superose 12 column equilibrated in PB (Pharmacia LKB Biotechnology) at a flow-rate of 0.4 ml/min. Fractions containing ribosome inactivating protein activity (as measured by a rabbit reticulocyte protein synthesis assay, described above) (the first major peak) were pooled. At this stage, the  $\alpha\beta$  RIP was usually quite pure as identified by SDS-PAGE (see Laemmli (1970), *supra*). If necessary, further purification can be achieved by applying the protein to a Mono S 5/5 column (Pharmacia LKB Biotechnology) equilibrated with PB and eluted at 1 ml/min with 0 to 50 mM sodium chloride in PB over 5 min, then 50 to 200 mM sodium chloride in PB over 25 min.

Results from a typical purification are presented in Table 5. The effect of purified maize  $\alpha\beta$  RIP on mammalian protein synthesis is shown in Figure 2.

Table 5

## Purification of maize RIP from mature kernels

Step	Protein (mg)	Total units <sup>1</sup> x 10 <sup>6</sup>	Yield (%)	Fold Purification	IC <sub>50</sub> (ng/ml)
Crude extract	6816	384	100	1.0	323
85% Ammonium sulfate	1010	115	30	2.0	161
post-DE52 treatment	428	144	38	5.9	54
Mono S 10/10 pool	10.2	58	15	102	3.2
Superose 12 pool	1.8	33	8.6	327	0.99
Mono S 5/5 pool	1.32	32.4	8.4	436	0.74

1. One unit of activity is the amount of protein required to produce 50% inhibition in the rabbit reticulocyte lysate protein synthesis assay.

## A. Rabbit Reticulocyte Cell-Free Protein Synthesis Assay

The inhibitory activity of the maize  $\alpha\beta$  RIP toward mammalian protein synthesis was measured in a

rabbit reticulocyte lysate system following the procedures of Pelham and Jackson (see (1976), Eur. J. Biochem., 67:247-256).

A mix of the following reagents was prepared (2.5 milliliter (ml) total volume): 125 microliter ( $\mu$ l) 200 mM Tris-HCl, pH 7.6 + 40 mM magnesium acetate + 1.6 M potassium chloride; 12.5  $\mu$ l 3 mM hemin hydrochloride in 50 percent ethylene glycol; 1.0 ml untreated rabbit reticulocyte lysate (Promega, Madison, WI); 1.0 ml H<sub>2</sub>O; 62.5  $\mu$ l amino acid mix; 125  $\mu$ l 20 mM ATP + 4 mM GTP; 125  $\mu$ l 200 mM creatine phosphate; 50  $\mu$ l 2.5 mg/ml creatine phosphokinase in 50 percent ethylene glycol. The amino acid mix contained 50  $\mu$ M of each amino acid except glycine (100  $\mu$ M), arginine, isoleucine, methionine and tryptophan (10  $\mu$ M each) and contained no leucine. All stock solutions were previously adjusted to pH 7.5 prior to addition.

Five microliters (5  $\mu$ l) of appropriate dilutions of samples to be assayed were placed in the wells of a 96-well plate and 50  $\mu$ l of the mix added. After 10 minutes, 50 nanoCuries (nCi) <sup>14</sup>C-leucine in 10  $\mu$ l was added to each well. After a further 10 minutes (min.), the reaction was quenched with 10  $\mu$ l 1.5 M potassium hydroxide and incubated for 45 min. Twenty-five microliters (25  $\mu$ l) of each sample was then pipetted onto individual 2.1 cm Whatman 3 MM paper disks (Whatman, Clifton, NJ) and after drying for 2 to 3 min, the disks were washed successively by swirling in a flask with 250 ml 10 percent trichloroacetic acid, 250 ml 5 percent trichloroacetic acid (twice), 125 ml ethanol, 250 ml 1:1 ethanol/acetone, and 125 ml acetone. After drying, the filters were placed in vials with 10 ml scintillation cocktail and counted.

#### 20 B. Antisera and Western blot analysis:

The  $\alpha$  and  $\beta$  polypeptide bands were cut from 3 millimeter (mm) SDS-PAGE gels after brief staining with Coomassie blue and were electroeluted using an electroelution device (Bio-Rad, Richmond CA) according to the manufacturer's directions. The polypeptide preparations were then used to immunize rabbits to yield polyclonal anti-sera (prepared by RIBI Immunochem, Montana).

Western blots from Phastgels™ reagent (Pharmacia LKB Biotechnology) were performed by removing the gel from the plastic backing and then electroblotting onto Immobilon paper (Millipore Corporation, Bedford, MA). Blots were developed using the maize  $\alpha\beta$  RIP primary antiserum at 1:2000 dilution and alkaline phosphatase-conjugated goat anti-rabbit secondary antibody (Bio-Rad), according to the manufacturer's instructions.

#### Example 2: Isolation of Maize proRIP

The polyclonal antisera against the  $\alpha$  and  $\beta$  fragments were used to identify a common 34 kD precursor polypeptide in crude extracts of maize kernels (maize proRIP). The presence of the maize proRIP was monitored during subsequent purification by Western blot analysis as set forth above. All steps of the purification were performed at 4 °C, except for HPLC which was performed at room temperature.

Two hundred fifty grams (250 g) of immature maize kernels were homogenized in 600 ml 25 mM sodium phosphate, pH 7.2 (PB) + 5  $\mu$ g/ml antipain. After the extract was strained through several layers of cheesecloth, the protein precipitating between 45 and 80 percent ammonium sulfate was collected and redissolved in 15 ml PB, then passed over a 2.5 x 15 cm Sephadex G-25 column (Pharmacia LKB) equilibrated in PB. Fractions containing protein were pooled and diluted to ~60 ml with water. The solution was applied to a Q-Sepharose (fast-flow) column packed in a 10/10 FPLC column (Pharmacia LKB Biotechnology) equilibrated with PB, and eluted with a 0 to 300 mM NaCl gradient at 2 ml/min over 75 min. Fractions containing the 34 kD precursor were pooled and concentrated by a Centriprep 10 device (Amicon) to 1.5 ml. This was diluted four-fold with water and applied to a Mono Q 5/5 column (Pharmacia LKB Biotechnology) equilibrated in PB. The column was eluted with a 0 to 250 mM NaCl gradient over 60 min. Fractions containing the 34 kD polypeptide were pooled, concentrated to 0.5 ml and applied to a Superose 12 column (Pharmacia LKB Biotechnology) equilibrated in PB. The major peak from this column contained the 34 kD maize RIP precursor and appropriate fractions were pooled and stored at -20 °C.

#### Example 3: PAGE Analysis of Maize $\alpha\beta$ RIP and proRIP

SDS-PAGE was performed with a Phastsystem™ reagent (Pharmacia LKB Biotechnology) using 20 percent Phastgels™ reagent and following the manufacturer's instructions. Native PAGE was performed at pH 4.2 as described in the Phastsystem™ reagent application file no. 300, method 1 (Pharmacia LKB Biotechnology).

SDS-PAGE of highly purified, active maize  $\alpha\beta$  RIP exhibited two polypeptides an  $\alpha$  fragment (16.5 kD)

and a  $\beta$  fragment (8.5 kD) under both reducing and non-reducing conditions. A single band was seen in native PAGE analysis of purified, active maize  $\alpha\beta$  RIP. The minimal Mr value of the associated, native maize  $\alpha\beta$  RIP was therefore 25 kD.

By SDS-PAGE, highly purified maize proRIP migrated with a value of 34 kD.

5

#### Example 4: *In vitro* activation of Maize proRIP by Papain

A purified sample of proRIP was incubated at 0.5 mg/ml with papain, a plant thiol protease, at 0.01 mg/ml in sodium acetate buffer, pH 6 containing 2 mM dithiothreitol. After 1 to 2 hours at room temperature, the 34 kD proRIP was digested to a stable product exhibiting a polypeptide pattern almost identical to that of native, active maize  $\alpha\beta$  RIP. There was a concomitant increase in ribosome inactivating activity in the incubation; the undigested proRIP had no ribosome inactivating activity up to 2  $\mu$ g/ml, whereas papain-treated proRIP had an IC<sub>50</sub> of <80 ng/ml. In contrast trypsin had no effect on maize proRIP.

#### 15 Example 5: Chemically-determined amino acid sequences

##### A. N-Terminal Amino Acid Sequences of Maize $\alpha\beta$ RIP $\alpha$ fragment and $\beta$

A sample of maize  $\alpha\beta$  RIP was electrophoresed by the method of Laemmli (1970), *supra* in 1.5 mm thick gels and the gel electroblotted onto Immobilon PVDF paper (Millipore) using a Transphor™ apparatus (Pharmacia LKB Biotechnology). The paper was stained briefly with Coomassie blue and the bands corresponding to the 16.5 and 8.5 kD polypeptides were cut out. These were N-terminal sequenced directly from the PVDF paper using an 470A gas phase sequencer (Applied Biosystems, Foster City, CA). The following data was obtained (bracketed residues denote lower confidence assignments):

25 N-Terminal sequence of  $\alpha$  fragment:

30 K R I V P K I T E I F P V E D A N Y P V S A F I A [G] V X  
K D V I

An additional minor species (~20 percent of the total species) had an N-Terminal sequence of:

35 A Q T N K[L]I V P K

N-Terminal sequence of  $\beta$  fragment:

40 A A D P Q A D T K S X L V K L V V M S/C E G L X F N T V  
S

45

##### B. $\alpha$ fragment C-Terminal Amino Acid Sequence

The carboxy-terminal amino acid sequence of the  $\alpha$  maize  $\alpha\beta$  RIP  $\alpha$  fragment was determined using sequencing grade carboxypeptidase P from *Penicillium japonicum* (Boehringer Mannheim, Indianapolis, IN). A sample of 16.5 kD polypeptide was purified by reverse-phase HPLC using a Vydac 5 $\mu$  C4 4.6 x 30 mm RP column. The column was equilibrated with 0.1 percent trifluoroacetic acid (TFA), and eluted with 0 to 40 percent of 0.1 percent TFA + 80 percent acetonitrile over 8 min, then 40 to 60 percent of 0.1 percent TFA + 80 percent acetonitrile over 20 min. The  $\beta$  fragment eluted after 21.9 min. and the  $\alpha$  fragment eluted after 23.3 min.

55 A lyophilized sample of the  $\alpha$  fragment was dissolved in 20 mM sodium acetate, pH 5.8 + 4 M urea. The digestion mix contained the following in 0.1 ml: 1.6  $\mu$ g carboxypeptidase P, 66  $\mu$ g  $\beta$  fragment, 0.12 M sodium acetate pH 4.2, 0.8 M urea. After 1, 5, 15, 60, 120 and 480 min, duplicate 8  $\mu$ l aliquots from the digestion were added to 8  $\mu$ l 0.4 M sodium borate, pH 10.5 and frozen on dry ice.

Amino acid analysis was performed essentially as described by Jones (1986), In: Methods of Protein Microcharacterization (ed.) J.E. Shively. The following sequence is obtained:  $\text{NH}_2\text{-Asp-Leu-Ala-(Lys)}_n\text{-COOH}$ , where  $n = 2\text{-}4$ . This was the carboxy terminus of the  $\alpha$  polypeptide, therefore this and the N-terminus sequence of the  $\beta$  fragment define the linker region contained in the precursor (see amino acid sequence derived from cDNA in Table 1).

#### C. N-Terminal Amino Acid Sequence of Maize proRIP

No N-Terminal sequence data was obtained from a sample of the 34 kD maize proRIP indicating that this polypeptide is N-terminal blocked.

#### Example 6: Isolation and Characterization of cDNA for Maize proRIP

##### A. Isolation

Immature kernels from field grown Pioneer hybrid 3737 were harvested, shelled from the cob, and stored at  $-20^\circ\text{C}$ . Ten grams (10 g) of kernels were frozen in liquid nitrogen for several minutes then ground to a powder in a Waring blender. The powder was suspended in 20 ml of ice cold TENS buffer (10 mM Tris pH 7.4, 1 mM EDTA, 0.5 percent SDS, 0.3 M NaCl) and extracted immediately with an equal volume of phenol-chloroform-isoamyl alcohol (25:24:1) saturated with TENS buffer. The aqueous phase was collected and extracted three more times with fresh phenol mixtures.

Nucleic acids were precipitated from the aqueous phase by adjusting it to 0.3 M sodium acetate pH 5.5 and adding 2.5 volumes of 100 percent ethanol. Nucleic acids were collected by centrifugation and suspended directly in 1 ml phenol-chloroform-isoamyl alcohol plus 1 ml TENS and extracted by vortexing. The nucleic acid was precipitated from the aqueous phase by ethanol precipitation as above. The precipitate was collected by centrifugation and resuspended in TE buffer (10 mM Tris pH 7.4, 1 mM EDTA). Single strand nucleic acid was precipitated by adjusting the solution to 2M LiCl and incubating for 4 to 12 hours at  $4^\circ\text{C}$ . Centrifugation yielded a pellet which consisted of between 2.2 to 2.5 mg of total RNA.

Poly(A)-containing RNA was enriched from the total RNA sample by using Hybond mAPTM mRNA purification affinity paper (Amersham Corporation, Arlington Heights IL). The supplier's protocol was followed. Typically 5 to 10  $\mu\text{g}$  of poly(A) enriched RNA were recovered per milligram of total RNA.

Five micrograms (5  $\mu\text{g}$ ) of poly(A) enriched RNA were converted into double stranded cDNA using a cDNA Synthesis<sup>TM</sup> kit (Pharmacia LKB Biotechnology). The cDNA was ligated into the cloning vector Lambda gt11 (Stratagene Inc., La Jolla CA) following the supplier's instructions. Packaging of the ligated vector-insert mixture was done with the Gigapack plus packaging extract (Stratagene, Inc.) again following the supplier's protocol.

The PicoBlue Immunodetection<sup>TM</sup> kit (Stratagene, Inc.) was used to screen the Lambda gt11 maize kernel cDNA library using rabbit polyclonal antisera raised against the maize proRIP, as described above.

Positive clones were purified to homogeneity and the cDNA inserts characterized by *Eco* RI restriction enzyme analysis. One of the largest *Eco* RI generated cDNA inserts (about 1,100 bp) was ligated into the *Eco* RI site of plasmid pUC19 (Bethesda Research Labs, Gaithersburg, MD). Clones carrying the proRIP cDNA insert in both orientations were identified by restriction digestion and used for large scale plasmid purification.

##### B. Sequencing the maize proRIP cDNA

The nucleotide sequence of the proRIP cDNA (set forth in Table 1) was determined by dideoxy chain termination sequencing using the Sequenase<sup>TM</sup> DNA sequencing kit (United States Biochemical Corp., Cleveland Ohio). The double stranded pUC19-RIP was used as template following the manufacturer's instructions. The first round of sequencing was initiated by the M13/pUC forward sequencing primer (Bethesda Research Labs). Subsequent primers were derived from the sequenced maize proRIP cDNA. Both strands of the cDNA were fully sequenced at least once.

The open reading frame encoding the  $\alpha\beta$  RIP protein was confirmed by comparing the cDNA deduced amino acid sequence (set forth in Table 1) to the chemically determined protein sequence data.

#### Example 7: Expression of Maize proRIP and Derivatives in *Escherichia coli*

Various genetic derivatives of maize proRIP may be expressed in *E.coli* and tested for ribosome

inactivating activity. A summary of several constructions and their properties is given below.

A. R34 (set forth in Table 6 below) represents the intact recombinant proRIP gene which encodes a protein of Mr 33,327 and as expected is not a potent inhibitor of protein synthesis. Upon papain treatment it is processed into two associated polypeptides (of approximately 17 + 9 kD) by SDS Phastgel™ analysis) with very potent ribosome inactivating activity. N34 represents the native proRIP as isolated from nature.

Table 6

**R34 DNA Sequence: The Maize ProRIP DNA Engineered for Expression in *Escherichia coli***

```


1  GCTTAATTAA TTAAGCTTAA AAGGAGGAAA AAAATTATCG CCGAGATAAC CTTAGAGCCG
61 AGTGATCTTA TGGCGCAAAC AAACAAAAGA ATAGTGCCAA AGTTCAGTGA AATCTTCCCC
121 GTGGAGGACG CGAACTACCC TTACAGCGCC TTCATCGCGT CGGTCCGGAA AGACGTGATC
181 AAACACTGCA CCGACCATAA AGGGATCTTC CAGCCCGTGC TGCCACCGGA GAAGAAGGTC
241 CCGGAGCTAT GGTTCACAC AGAGCTCAA ACTAGGACCA GCTCCATCAC GCTCGCCATA
301 CGCATGGACA ACCTGTACCT CGTGGGCTTC AGGACCCCGG GCGGGGTGTG GTGGGAGTTC
361 GGCAAGGACG GCGACACCCA CCTCCTCGGC GACAACCCCA GGTGGCTCGG CTTGGCGGGC
421 AGGTACCAGG ACCTCATCGG CAACAAGGGT CTGGAGACCG TCACCATGGG CCGCGCCGAA
481 ATGACCAGGG CCGTCAACGA CCTGGCGAAG AAGAAGAAGA TGGCGACACT GGAGGAGGAG
541 GAGGTGAAGA TGCAGATGCA GATGCCGGAG GCCGCTGATC TGGCGGCGGC GGCAGCGGCT
601 GACCCACAGG CCGACACGAA GAGCAAGCTG GTGAAGCTGG TGGTCATGGT GTGCGAGGGG
661 CTGCGGTTCA ACACCGTGTC CCGCACGGTG GACGCGGGGT TCAACAGCCA GCACGGGGTG
721 ACCTTGACCG TGACGACGGG GAAGCAGGTG CAGAAGTGGG ACAGGATCTC CAAGGCGGCC
781 TTCGAGTGGG CTGACCAACC CACCGCTGTG ATCCCGGACA TGCAGAAGCT TGGCATCAAG
841 GATAAGAAGC AAGCAGCGAG GATCGTTGGG CTCSTTAAGA ATCAAACCTAC TGCCGCTGCC
901 GCTACTGCTG CCAGTGCTGA CAACGACGAC GACGAGGCCT GATCAATGCA ACGACACATC
961 ATGATCTGCT GGTGCACTTA ATTACTATGT TCGTATACAA ATAAATACAC CCGGCGTACG
1021 CCGTGTTCTT TATATGGTCT AAAATGTAGC CAGTAAATTT TAACTACTT TCTCGTGGCG
1081 AATTCAGTGG CCGGCGTGGT ATATA

```

Expression of the recombinant maize proRIP in *E. coli* was accomplished by engineering the cDNA via PCR amplification. A 5' primer was synthesized which contained termination codons in all three reading frames to stop translation of vector-encoded proteins upstream of the maize proRIP cDNA. The primer also contained a Shine-Dalgarno sequence several base pairs upstream of an ATG start codon followed by 23 bases which were homologous to the maize proRIP cDNA. The 3' primer spans the 3' cDNA end-pUC19 junction (the primers were shown by the underlined regions set forth in Table 6). PCR amplification of the cDNA in pUC19 using the GeneAmp™ kit (Perkin Elmer-Cetus, Norwalk, CT) yielded a predominant amplification product of approximately 1100 base pairs, as expected.

The engineered, amplified product, was purified from an agarose gel and ligated into the filled-in *Hind* III

site of the expression vector pGEMEX-1 (Promega Corp., Madison, WI) to give plasmid pGR, set forth below.



This was transformed into *E. coli* DH5 $\alpha$  (Bethesda Research Labs). Plasmids containing the maize proRIP cDNA were isolated by colony hybridization (see Sambrook *et al.*, *supra*) with a 5' maize proRIP cDNA probe and characterized. Those containing the cDNA probe for the maize proRIP in the correct orientation were tested for expression. Plasmids were transformed into competent *E. coli* JM109(DE3) (Promega Corp., Madison, WI), transformed cells were grown in 15 ml cultures under ampicillin selection to an optical density at 600 nm of 0.4 to 1.0. Isopropylthio- $\beta$ -galactoside (IPTG) was added to 1.3 mM to induce the production of recombinant proRIP and the cultures were grown an additional 4 hours at 37 $^{\circ}$  C. The cells were collected by centrifugation and stored as a pellet at -20 $^{\circ}$  C.

The protein produced from the maize proRIP cDNA was analyzed by lysing the induced cells in TE containing 1 mg/ml lysozyme 37 $^{\circ}$  C for 15 min. The lysate was fractionated into a crude supernatant and pellet by microcentrifugation. The fractions were analyzed by SDS-PAGE using 20 percent Phastgels<sup>TM</sup> reagent (Pharmacia LKB Biotechnology). Coomassie blue staining and Western blot analysis of the gels with anti-maize proRIP sera identified a 34 kD band which was greatly increased upon induction of the cells with IPTG. Cells not carrying the plasmid or containing the plasmid with the maize proRIP cDNA in the inverted orientation did not contain this 34 kD immuno-reactive band. The majority of the recombinant maize proRIP was contained in the cellular pellet suggesting the material was insoluble under these conditions.


To test if R34 could acquire the folding pattern of N34 the pellet fraction of an induced culture was dissolved in 6M guanidine HCl and allowed to denature at room temperature for 3 hours. The material was then diluted 200-fold into ice cold TE and incubated at 4 $^{\circ}$  C overnight to allow refolding of the denatured R34. The diluted material was then concentrated by a Centricon 10 device (Amicon). To test whether refolded R34 could undergo the correct proteolytic processing to the fragmented form of the maize proRIP, the material was treated with 10  $\mu$ g/ml papain for various times, and samples were analyzed by SDS-Phastgel and Western blot analysis. The R34 material was processed to a stable mixture of two immuno-reactive bands which comigrate with N34 papain-processed material indicating the correct proteolytic processing had occurred.

In an effort to simplify purification of the R34 polypeptide from induced lysates, the gene 10 coding region of the pGEMEX-1 vector was removed by cutting the maize proRIP gene-containing plasmid (pGR) with *Xba* I and gel purifying the vector/proRIP DNA away from the gene 10 encoding DNA. Recircularization of pGR, now minus the gene 10 coding region, resulted in a plasmid called pGR1 set forth below.



5

10



15 The plasmid pGR1 was transformed into JM109(DE3) cells and tested for production of R34 following induction with IPTG. As with pGR, large amounts of R34 were identified in cellular lysates both by Western blot and Coomassie blue staining. Unlike pGR, R34 produced from pGR1 was soluble and fractionated in the supernatant of lysed cells. This soluble material was treated with papain at 10  $\mu$ g/ml and the R34 produced from pGR1 was cleaved to products which comigrate with N34, papain-cleaved product. The

20 papain-treated material inhibited translation of reticulocyte lysates at significantly higher dilutions than the untreated material, indicating that the soluble R34 was processed to an active form.

B. R34-DL represents the proRIP without the linker. The sequences encoding  $\alpha$  and  $\beta$  were joined directly without intervening linker DNA, i.e., nucleotides A-520 to A-594 are deleted. The R34-DL gene encoded a 30.6 kD protein which was a potent inhibitor of protein synthesis. Treatment of R34-DL with

25 papain resulted in a 28 kD polypeptide with increased ribosome inactivating activity over the untreated molecule.

Confirmation that removal of the linker from maize proRIP activated the molecule was obtained independently through genetic engineering. The 75 bp linker encoding region of R34 (A-520 to A-594 inclusive, Table 6) was deleted using PCR amplification. The new construction R34-DL joined directly, in

30 frame, the DNA encoding both the  $\alpha$  and  $\beta$  fragments.

In the pGEMEX-1 system the R34-DL gene directed the synthesis of a polypeptide approximately 30.6 kD, which was recognized by antisera specific for the maize proRIP. At high dilution, *E.coli* lysates containing R34-DL protein were potent inhibitors of protein synthesis in rabbit reticulocyte lysates, in marked contrast to *E. coli* lysates containing the R34 polypeptide.

35 These genetic deletion data confirm that removal of the linker served to activate the R34 (proRIP) molecule. This experiment also demonstrated that covalent linkage of the  $\alpha$  and the  $\beta$  polypeptide fragments resulted in an active  $\alpha\beta$  RIP. The maize proRIP did not require a break in the polypeptide backbone for enzymatic activity, removal of the linker region was sufficient to confer potent ribosome inactivating activity.

40 In addition, when R34-DL lysates were treated with papain a slight decrease in the molecular weight of R34-DL protein is noted (from 30.6 kD to approximately 28 kD). The R34-DL polypeptide remained intact, that is, it was not cleaved to the characteristic maize  $\alpha\beta$  RIP  $\alpha$  and  $\beta$  fragments. Associated with this small change in molecular weight was an increase in protein synthesis inhibition in the *E.coli* lysates. These data indicated that in bacterial lysates removal of the linker region activated the ribosomal inactivating activity of

45 the protein at least 250-fold, but that additional processing from the ends of the protein increased the activity.

Another genetic construction was made using PCR technology to remove the leader region from R34-DL. The new construction called R30-DL (nucleotides C-40 to C-84 and A-520 to A-594 inclusive are deleted) encoded a protein (approximately 29.5 kD) which was slightly smaller than R34-DL. *E.coli* lysates

50 containing R30-DL appeared to be even more potent inhibitors of protein synthesis than R34-DL lysates. Papain treatment of R30-DL containing lysates further enhanced protein synthesis inhibiting activity. Following this treatment the R30-DL protein underwent a slight decrease in molecular weight representing processing at the ends of the polypeptide..

55 Examp1 8:

A segment of the R30-DL gene was deleted which encodes several acidic residues at the carboxy terminus of the protein. The deletion was accomplished using the PCR engineering methods.

A thermocycler (Perkin-Elmer Cetus, Norwalk, CT) was used for the indicated constructions. A typical run was done with a one minute denaturation step, 2 minute annealing, and a 3 minute extension step. Temperatures used were 94 °C, 37 °C or 50 °C, and 72 °C respectively. Following 25 cycles the reaction was held at 72 °C for 7 minutes for extension of unfinished products.

Amplification engineering reactions were done in four separate tubes of 100 µL each. The tubes were combined following amplification. Normally 100 ng of template was included in each tube. DNA primers were synthesized on a PCR Mate or 380A DNA synthesizer (Applied Biosystems) and were purified on acrylamide gels. Fifty (50) pmol of each primer were included in each reaction. The reaction conditions for the amplification were those recommended by Perkin Elmer Cetus (10 mM Tris-HCl pH8.3, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.001 percent gelatin, 200 µM dNTPs and 2.5 units of *Taq* DNA polymerase or AmpliTaq™ thermostable DNA polymerase).

Several methods of genetic engineering were employed to produce the genetic derivatives described below. The standard methods of DNA purification, restriction enzyme digestion, agarose gel analysis, DNA fragment isolation, ligation and transformation were as described (see Molecular Cloning: A Laboratory Manual, *supra* and Current Protocols in Molecular Biology, (1987), *supra*).

Enzymes used for the engineering were from one of three manufacturers (Pharmacia LKB Biotechnology; Bethesda Research Labs; or New England Biolabs, Beverly, MA). Buffers and protocols used were provided by the manufacturer.

Using PCR introduced engineering, a modified RIP fragment is amplified from an RIP plasmid template, purified, then used to replace the unmodified region in the RIP gene. All fragment replacements were done in RIP genes already inserted into a pGEMEX expression plasmid.

A pGEMEX plasmid containing R30-DL, which had the 3' half of the gene removed by *Nco* I and *Stu* I digestion followed by gel purification was used as the template for PCR. The 3' half of the RIP gene was replaced with the PCR modified fragment described below.

A 3' PCR primer was synthesized which encoded the 7 amino acid deletion near the carboxy terminus and introduced a new unique *Bam* HI site. The 5' primer directed the deletion of the  $\alpha\beta$  linker and included a *Nco* I site. The sequences of these primers is given in Table 7. The primers were used to amplify a modified DNA fragment from a pGEMEX R34-DL template. The amplified fragment was phenol extracted and ethanol precipitated. The insert DNA was cut with *Nco* I and ligated into the pGEMEX-R30-DL vector.

Table 7

### Polymerase Chain Reaction primers for RDT.

Table 7: Polymerase Chain Reaction primers for RDT.

5' Primer

5' ACC CTC ACC ATG GGC GGC GGC GAA ATG ACC AGG GGC CTC AAC GAC CTC GGC AAG  
AAG AAG AAG GGC GGT GAC CCA CAG GGC GAC ACG AAG AGC 3'

3' Primer

5' GGC ATG CAG CAG TAG GGC GAG GGC CAG TAG 3'

The new RIP gene derivative is called RDT and encodes a protein of predicted a 28,233 Daltons and pI ~9.5. The RDT gene encodes a protein with a truncated leader, deleted linker and truncated carboxy terminus.

The DNA sequence for RDT is shown in Table 8.

Table 8

Nucleotide sequence and deduced amino acid sequence of  
the maize RIP derivative RDT

Table 8: Nucleotide sequence and deduced amino acid  
sequence of the maize RIP derivative RDT.

15	1	TCCTCTAGATGGGGCTAATTAACTAAGCTTAAAGGAGCAAAAAAATT	ATG
	2		Met
	54	AAA AGA ATA GTG CCA AAG TTC ACT GAA ATC TTC CCC GTG GAG GAC GCG AAC	
	55	Lys Arg Ile Val Pro Lys Phe Thr Glu Ile Phe Pro Val Glu Asp Ala Asn	
	105	TAC GGT TAC AGC GGC TTC ATG GCG TCG GTC CCG AAA GAC GTG ATC AAA CAC	
	106	Tyr Phe Tyr Ser Ala Phe Ile Ala Ser Val Arg Lys Asp Val Ile Lys His	
20	156	TGC ACC GAC CAT AAA GGG ATC TTC CAG CCC GTG CTG CCA CCG GAG AAG AAG	
	157	Cys Thr Asp His Lys Gly Ile Phe Gln Pro Val Leu Pro Pro Glu Lys Lys	
	207	GTG CCC GAG CTA TGG TTC TAC ACA GAG CTC AAA ACT AGG ACC AGC TCC ATC	
	208	Val Pro Glu Leu Trp Phe Tyr Thr Glu Leu Lys Thr Arg Thr Ser Ser Ile	
25	258	AGC CTC GGC ATA GCG ATG GAC AAC CTC TAC CTC GTG GGC TTC AGG ACC CCG	
	259	Thr Leu Ala Ile Arg Met Asp Asn Leu Tyr Leu Val Gly Phe Arg Thr Pro	
	309	GGC GGC GTG TGG TGG GAG TTC GGC AAG GAC GGC GAC ACC CAC CTC CTC GGC	
	310	Gly Gly Val Trp Trp Glu Phe Gly Lys Asp Gly Asp Thr His Leu Leu Gly	
	360	GAC AAC CCC AGG TGG CTC GGC TTC GGC GGC AGG TAC CAG GAC CTC ATC GGC	
	361	Asp Asn Pro Arg Trp Leu Gly Phe Gly Gly Arg Tyr Gln Asp Leu Ile Gly	
30	411	AAC AAG GGT CTG GAG ACC GTC ACC ATG GGC CCC GGC GAA ATG ACC AGC GGC	
	412	Asn Lys Gly Leu Glu Thr Val Thr Met Gly Arg Ala Glu Met Thr Arg Ala	
	462	GTC AAC GAC CTC GCG AAG AAG AAG AAG GCG GCT GAC CCA CAG GGC GAC ACC	
	463	Val Asn Asp Leu Ala Lys Lys Lys Lys Ala Ala Asp Pro Gln Ala Asp Thr	
35	513	AAG ACC AAG CTC GTG AAG CTC GTG GTC ATG GTG TGC GAG GGC CTC CGG TTC	
	514	Lys Ser Lys Leu Val Lys Leu Val Val Met Val Cys Glu Gly Leu Arg Phe	
	564	AAC ACC GTG TCC GCG ACC GTG GAC GCG GGC TTC AAC AGC CAG CAC GGC GTG	
	565	Asn Thr Val Ser Arg Thr Val Asp Ala Gly Phe Asn Ser Gln His Gly Val	
	615	ACC TTC ACC GTG ACC CAG GCG AAG CAG GTG CAG AAG TCG GAC AGG ATC TTC	
	616	Thr Leu Thr Val Thr Gln Gly Lys Gln Val Gln Lys Trp Asp Arg Ile Ser	
40	666	AAG GCG GCG TTC GAG TGG GGT GAC CAC CCC ACC GGT GTG ATC CCC GAC ATG	
	667	Lys Ala Ala Phe Glu Trp Ala Asp His Pro Thr Ala Val Ile Pro Asp Met	
	717	TAG AAG GTT GCG ATC AAG CAT AAG AAC GAA GCA GCG AGG ATC GTT GCG CTC	
	718	Tin Lys Leu Gly Ile Lys Asp Lys Asn Glu Ala Ala Arg Ile Val Ala Leu	
45	769	GTG AAG AAT GAA ACT ACT GCG GGT GCG GGT ACT GGT GGA TCG GCG TGA TGA	
	770	Val Lys Asn Glu Thr Thr Ala Ala Ala Ala Thr Ala Gly Ser Ala End	
	819	ATGCAACGACACATCATCATCTGCTGCTGCACTTAATTACTATGCTGCTATACAAATAATACACCC	
	869	TCGTAAGCGGTGTCTCTTATGCTGCTTAAATGTAGCGCAATTAATTTAAAGTACTTTCTGCTGCG	
50	919	TAATTCAGTGGCGGCATGCTATATA	

The RDT gene, expressed in *E.coli* using the pGEMEX system described above, was purified from bacterial lysates to apparent homogeneity. RDT protein appears to be a more potent inhibitor of protein synthesis than R30-DL. Using the reticulocyte lysate protein synthesis assay, purified RDT has a IC<sub>50</sub> value of 1 ng/ml.

Example 9:

RDT was further engineered to produce another gene called RDT-NP. This construction differs from RDT in having two unique restriction sites engineered into the gene. The sites were introduced using PCR methods described in Example 8. The PCR primer was designed such that it included the desired change and a unique restriction site in the maize RIP DNA sequence. A 99 bp primer which introduced the *Not* I and *Pst* I sites at the 3' end of the primer and had to be built back to the unique *Nco* I site for cloning purposes. The primers for the amplification are shown in Table 9.

Table 9

Polymerase chain reaction primers for RDT-NP.

Table 9: Polymerase chain reaction primers for RDT-NP.

5' Primer

5' ACC GTC ACC ATG GCG GCG GCG TCA ATG ACC AGG GCG GTC AAC GAG GTC GCG AAG  
TAG AAG AAG GCG GCG GCG TTT TTT TTT GAG TCA GAG GCG GAG ACC AAG

3' Primer

3' GAT GCG GCG GAG TCA ATT GCG

The restriction sites (*Not* I and *Pst* I) correspond to the site of the alpha/beta linker insertion in the RIP polypeptide. RDT-NP allows DNA segments encoding various polypeptide linkers to be inserted into the gene and tested for their ability to create an inactive, yet protease activatable RIP. The sequence for RDT-NP is shown in Table 10.

Table 10

Predicted nucleotide sequence and deduced amino acid  
sequence of the maize RIP derivative RDT-NP

Table 10: Predicted nucleotide sequence and deduced amino acid  
sequence of the maize RIP derivative RDT-NP

5	1	TTCTCTAGATGGGGCTAATTAAATTAAGCTTAAAGGAGGAAAAAAATG ATG AAA
	1	Met Lys>
10	57	AGA ATA GTG GCA AAG TTC ACT GAA ATC TTC GCG GTG GAG GAG GCG
	3	Arg Ile Val Pro Lys Phe Thr Glu Ile Phe Pro Val Glu Asp Ala>
15	102	AAC TAC GGT TAC AGC GCG TTC ATC GCG TCG GTG GCG AAA GAC GTG
	108	Asn Tyr Pro Tyr Ser Ala Phe Ile Ala Ser Val Arg Lys Asp Val>
	147	ATC AAA CAC TCG ACC GAC CAT AAA GCG ATC TTC CAG GCG GTG CTG
	153	Ile Lys His Cys Thr Asp His Lys Gly Ile Phe Gln Pro Val Leu>
20	192	GCA GCG GAG AAG AAG GTC GCG GAG CTA TGG TTC TAC ACA GAG CTC
	198	Pro Pro Glu Lys Lys Val Pro Glu Leu Trp Phe Tyr Thr Glu Leu>
	237	AAA ACT AGG ACC ACC TCC ATC ACC CTC GCG ATA GCG ATG GAC AAC
	243	Lys Thr Arg Thr Ser Ser Ile Thr Leu Ala Ile Arg Met Asp Asn>
	282	TTC TAC CTC GTG GCG TTC AGG ACC GCG GCG GCG GTG TCG TCG GAG
	288	Leu Tyr Leu Val Gly Phe Arg Thr Pro Gly Gly Val Trp Trp Glu>
25	327	TTC GCG AAG GAC GCG GAC ACC GAC CTC CTC GCG GAC AAC GCG AGG
	333	Phe Gly Lys Asp Gly Asp Thr His Leu Leu Gly Asp Asn Pro Arg>
	372	TGG CTC GCG TTC GCG GCG AGG TAC CAG GAC CTC ATC GCG AAC AAG
	378	Trp Leu Gly Phe Gly Gly Arg Tyr Gln Asp Leu Ile Gly Asn Lys>
30	417	GCT CTC GAG ACC GTC ACC ATG GCG GCG GCG GAA ATG ACC AGG GCG
	423	Gly Leu Glu Thr Val Thr Met Gly Arg Ala Glu Met Thr Arg Ala>
		Not1 Not1
	462	TTC AAC GAC CTC GCG AAG AAG AAG AAG GCG GCG GCG GGT GCA GAC
	468	Ile Asn Asp Leu Ala Lys Lys Lys Lys Ala Ala Ala Ala Asp>
35	507	GA GAG GCG GAG GCG AAG GCG GCG GCG GCG GCG GCG GCG GCG
	513	Pro Glu Ala Asp Thr Glu Ser Lys Leu Ala Ala Lys Ser Val Val Met>
	552	TTC TCG GAG GCG CTC GCG TTC GAG ACC GTC TCG GCG ACC GTC GAG
	558	Ala Cys Glu Gly Leu Arg Phe Asn Thr Val Ser Arg Ile Val Asp>
	597	TC GCG TTC AAC ACC GAG GAG GCG GCG GCG GCG GCG GCG GCG
	603	Ala Gly Phe Asn Ser Glu His Gly Val Thr Leu Thr Val Thr Glu>
40	642	CG AAG GAG GTG GAG AAG TCG GAG AGG ATC TCG AAG GCG GCG TTC
	648	Gly Lys Glu Val Glu Lys Pro Asp Arg Ile Ser Lys Ala Ala Phe>
	687	GAG TGG GGT GAG GAG TCG ACC GGT GTG ATC GCG GAG ATG GAG AAG
	693	Ile Trp Ala Asp His Pro Thr Ala Val Ile Pro Asp Met Glu Lys>
45	732	TTC GCG ATC AAG GAT AAG AAC GAA GCA GCG AGG ATC GTC GCG CTC
	738	Leu Gly Ile Lys Asp Lys Asn Glu Ala Ala Arg Ile Val Ala Leu>
	777	TTC AAG AAT GAA ACT ACT GCG GGT GCG GGT ACT GGT GGA TCG GCG
	783	Val Lys Asn Glu Thr Thr Ala Ala Ala Ala Thr Ala Gly Ser Ala>
	822	TCATCAATGCAACGACACATCATCATTTCTGCTGCACTTAATTACTATGTTCTATACA
		End
50	862	TATAAATACACCGGGGTACCGGGTCTTCTGCTTATATGGTCTAAAAATGAGCGAATAAT
	868	TAAAGTACTTCTGCTGCGCAATTCTCTGCGGGCATGCTATATA

The RDT-NP polypeptide had a predicted molecular weight of 28,446 Daltons and pI of 9.5. Crude  
lysates of *E.coli* expressing RDT-NP from a pGEMEX vector are potent inhibitors of eukaryotic protein  
synthesis.

Exempl 10:

## EP 0 466 222 A1

To create an RIP molecule which would bind to immunoglobulin IgG, a single Antibody Binding Region (ABR) domain from the *Staphylococcus aureus* antibody binding Protein A was subcloned from the plasmid pRIT5 (Pharmacia LKB Biotechnology) using PCR techniques. The antibody binding domain of protein A (ABR-A) was PCR engineered to have a *Bam* HI site at its 5' end and a *Bgl* II site at its 3' end. This allowed  
5 insertion of the ABR-A domain into the RDT *Bam* HI site while retaining the unique *Bam* HI site. The sequence of RDT-A is shown in Table 11.

10

15

20

25

30

35

40

45

50

55

Table 11

Predicted nucleotide sequence and deduced amino acid sequence of RDT-A.

5

10

15

20

25

30

35

40

45

50

```

1      TGGCTGTAGATGGGGCTAATTAAATTAGCTTAAAGGAGGAAAAAAATT ATG AAA
2      Met Lys>
3
4
5
6
7
8
9
10     AGA ATA GTG CCA AAG TTC ACC GAA ATC TTC GCG GTG GAG GAC GCG
11     Arg Ile Val Phe Lys Phe Thr Glu Ile Phe Pro Val Glu Asp Ala>
12
13     AAC TAC GGT TAC AGC GCG TTC ATC GCG TCG GTC GCG AAA GAC GTG
14     Asn Tyr Phe Tyr Ser Ala Phe Ile Ala Ser Val Arg Lys Asp Val>
15
16     ATC AAA CAC TCG ACC GAC GAT AAA GGG ATC TTC CAG CCG GTG CTG
17     Ile Lys His Cys Thr Asp His Lys Gly Ile Phe Gln Pro Val Leu>
18
19
20     CCA CCG GAG AAG AAG GTC CCG GAG CTA TGG TTC TAC ACA GAG CTC
21     Pro Pro Glu Lys Lys Val Phe Glu Leu Trp Phe Tyr Thr Glu Leu>
22
23     AAA ACT AGG ACC ACC TCG ACC ACC CTC GCG ATA GCG ATG GAC AAC
24     Lys Thr Arg Thr Ser Ser Ile Thr Leu Ala Ile Arg Met Asp Asn>
25
26     CTC TAC CTC GTG GCG TTC AGG ACC GCG GCG GGG GTG TCG TCG GAG
27     Leu Tyr Leu Val Gly Phe Arg Thr Pro Gly Gly Val Trp Trp Glu>
28
29
30     TTC GGC AAG GAC GCG GAC ACC CAC CTC CTC GCG GAC AAC CCC AGG
31     Phe Gly Lys Asp Gly Asp Thr His Leu Leu Gly Asp Asn Pro Arg>
32
33     TCG CTC GCG TTC GCG GCG AGG TAC CAG GAC CTC ATC GCG AAC AAG
34     Trp Leu Gly Phe Gly Gly Arg Tyr Gln Asp Leu Ile Gly Asn Lys>
35
36
37     GGT CTG GAG ACC GTC ACC ATG GCG GCG GCG CAA ATG ACC AGG GCG
38     Gly Leu Glu Thr Val Thr Met Gly Arg Ala Glu Met Thr Arg Ala>
39
40     CTC AAC GAC CTC GCG AAG AAG AAG AAG GCG GGT GAC CCA CAG GCG
41     Val Asn Asp Leu Ala Lys Lys Lys Lys Ala Ala Asp Pro Gln Ala>
42
43     TAC ACC AAG ACC AAG CTC CTC AAG CTC CTC CTC ATG CTC TCG GAG
44     Thr Thr Lys Thr Lys Leu Val Lys Leu Val Val Met Val Cys Glu>
45
46
47     TCG CTC GCG TTC AAG ACC CTC TCG GCG ACC CTC GAG GCG GCG TTC
48     Tyr Leu Arg Phe Asn Thr Thr Ser Arg Thr Val Asp Ala Gly Phe>
49
50     AAC ACC GAG CAG GCG GCG ACC TTC ACC CTC ACC CAG GCG AAG CAG
51     Asn Ser Gln His Gly Val Thr Leu Thr Val Thr Gln Gly Lys Gln>
52
53     CTG CAG AAG TCG GAC ACC ATC TCG AAG GCG GCG TTC CAG TCG GGT
54     Val Gln Lys Trp Asp Arg Ile Ser Lys Ala Ala Phe Glu Trp Ala>
55
56     GAC CAC CCG ACC GAT GTC ACC GCG GAC ATG CAG AAG CTT GCG ATC
57     Asp His Phe Thr Ala Val Ile Phe Asp Met Gln Lys Leu Gly Lys>
58
59     AAG GAT AAG AAC GAA GCA GCG AGG ATC GTC GCG CTC GTC AAG AAT
60     Lys Asp Lys Asn Glu Ala Ala Arg Ile Val Ala Leu Val Lys Asn>
61
62     GAA ACT ACT GCG GGT GCG GGT ACT GGT GCA TCG GGT GAT AAC AAT
63     Gln Thr Thr Ala Ala Ala Ala Thr Ala Gly Ser Ala Asp Asn Asn>
64
65     TTC AAC AAA GAA CAA CAA AAT GGT TTC TAC GAA ATC TTC AAT ATC
66     Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met>
67
68     GGT AAC TTA AAG GAA GAA CAA GCG AAT GGT TTC ATC CAA ACC TTA
69     Phe Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu>
70
71     AAA GAT GAC CCA AGC CAA AGT GGT AAC CTA TTC TCA CAA GGT AAA
72     Lys Asp Asp Pro Ser Gln Thr Ala Asn Leu Leu Ser Glu Ala Lys>
73
74     AAG TTA AAT CAA TGT CAA CAA TGT AAA GAT CCA TCG GCG TCAATCA
75     Lys Leu Asn Val Ser Gln Ala Phe Lys Asp Arg Ser Ala End<
76
77     TTTACGACACATATGATGATTTTCTTTCTTTTATTAATATATATTTTATACAAATATA
78
79     TTTACCGCGGATATCGGCTTTTCTTTTATATGCTTTTAAATGTATAGCGATAAATTTTAAA
80
81     TTAGCTTTTCTGTCGCAATTCTCTGCGGCGGATGCTATATA

```

RDT-A was expressed in *E.coli* cells using the pGEMEX system. The resulting polypeptide had a predicted molecular weight of 35,198 Daltons and pI of 9.2. It was recognized by antisera to both protein A and maize RIP indicating the chimeric nature of the protein. Crude lysates of bacteria expressing RDT-A had potent eukaryotic protein synthesis inhibition activity.

RDT-A was shown to bind specifically to IgG Sepharose (Pharmacia LKB Biotechnology) following the

manufacturer's instructions. Binding was best at pH 7.0. When washed at pH 5.0 the chimeric protein was released in small but detectable quantities from the resin. RDT alone does not bind to the gel.

#### Example 11:

5

To increase the binding ability of the RDT-A to IgG antibodies the Antibody Binding Domain from Streptococcal Group G protein G (ABR-G) was synthesized using oligonucleotides. The sequence synthesized was that of the naturally occurring sequence described by Guss *et al.* ((1986), *EMBO Journal*, 5:1567-1575). The only change was the addition of *Bam* HI and *Bgl* II sites at the 5' and 3' ends respectively of the synthetic DNA.

10

The ABR-G fragment was inserted into the *Bam* HI site of RDT-A. Two classes of clones have been studied. RDT-G-A contains a single ABR-G domain inserted in the correct orientation between the 3' end of RDT and the 5' end of ABR-A. A second class contains two properly oriented ABR-G domains. The predicted DNA sequences for the genes are shown in Tables 12 and 13.

15

Table 12

#### Predicted nucleotide and deduced amino acid sequence for RDT-G-A.

25	1	TCCTCTAGATCGGGCCTTAATTATTAAGCTTAAAGGAGGAAAAAATT ATG AAA
	2	Met Lys>
	3	ACA ATA GTG CCA AAG TTC ACT GAA ATC TTC CGC GTG CAG GAC GCG
	6	Arg Ile Val Pro Lys Phe Thr Glu Ile Phe Pro Val Glu Asp Ala>
	102	AAC TAC CTT TAC AGC GGC TTC ATC GCG TCG GTC CCG AAA GAC GTG
	15	Asn Tyr Pro Tyr Ser Ala Phe Ile Ala Ser Val Arg Lys Asp Val>
30	147	ATC AAA CAC TGC ACC GAC CAT AAA GCG ATC TTC CAG CCC GTG CTC
	33	Ile Lys His Cys Thr Asp His Lys Gly Ile Phe Gln Pro Val Leu>
	192	CCA CGC GAG AAG AAG GTC GCG GAG CTA TGG TTC TAC ACA GAG CTC
	48	Pro Pro Glu Lys Lys Val Pro Glu Leu Trp Phe Tyr Thr Glu Leu>
	237	AAA ACT AGG ACC AGC TCC ATC ACS CTC GGC ATA CCG ATG GAC AAC
35	63	Lys Thr Arg Thr Ser Ser Ile Thr Leu Ala Ile Arg Met Asp Asn>
	282	CTG TAC CTC GTG GGC TTC AGG ACC CCG GGC GCG GTG TGG TGG GAG
	78	Leu Tyr Leu Val Gly Phe Arg Thr Pro Gly Gly Val Trp Trp Glu>
	327	TTC GGC AAG GAC GGC GAC ACC CAC CTC CTC GGC GAC AAC CCC AGG
	93	Phe Gly Lys Asp Gly Asp Thr His Leu Leu Gly Asp Asn Pro Arg>
40	372	TGG CTC GGC TTC GGC GGC AGG TAC CAG GAC CTC ATC GGC AAC AAG
	108	Trp Leu Gly Phe Gly Gly Arg Tyr Gln Asp Leu Ile Gly Asn Lys>
	417	GGT CTG GAG ACC GTC ACC ATG GGC CCG GCG GAA ATG ACC AGG GCG
	123	Gly Leu Glu Thr Val Thr Met Gly Arg Ala Glu Met Thr Arg Ala>
	462	ATC AAC GAC CTG GCG AAG AAG AAG AAG GCG GGT GAC CCA CAG GCG
45	138	Val Asn Asp Leu Ala Lys Lys Lys Lys Ala Ala Asp Pro Gln Ala>
	507	GAC AGG AAG ACC AAG CTG CTG AAG CTG CTG CTG ATG GTG TCG GAG
	153	Asp Thr Lys Ser Lys Leu Val Lys Leu Val Val Met Val Cys Glu>
	552	TGG CTG GCG TTC AAG ACC GTC TCG GCG ACC GTC GAC GCG GCG TTC
	158	Ily Leu Arg Phe Asn Thr Val Ser Arg Thr Val Asp Ala Gly Phe>
50	597	GAC AGG GAC GAC GCG CTG ACC TTC ACC GTC ACC GAG GCG AAG CAG
	193	Asn Ser Gln His Gly Val Thr Leu Thr Val Thr Gln Gly Lys Glu>
	642	TTC CAG AAG TGG GAC AGG ATC TCT AAG GCG GCG TTC CAG TCG GGT
	198	Val Gln Lys Trp Asp Arg Ile Ser Lys Ala Ala Phe Glu Trp Ala>
	687	GAC CAG GCG ACC GGT GTC ATC GCG GAC ATG CAG AAG GTT GCG ATC
55	213	Val His Pro Thr Ala Val Leu Pro Asp Met Gln Lys Leu Gly Ile>
	732	AAG GAT AAG AAC GAA GCA GCG AGG ATC GTT GCG CTC GTT AAG AAT
	228	Lys Asp Lys Asn Glu Ala Ala Arg Ile Val Ala Leu Val Lys Asn>



Table 12 (cont.)

5

10

15

20

25

30

35

40

45

50

55

```

777   CAA ACT ACT GGC GGT GGC GGT ACT GGT GGA TCC AAA CCA GAA GGG
243   Gln Thr Thr Ala Ala Ala Ala Thr Ala Gly Ser Lys Pro Glu Val>

922   ATC GAT GCG TGT GAA TTA ACA CCA GCG GTG ACA ACT TAC AAA GTT
258   Ile Asp Ala Ser Glu Leu Thr Pro Ala Val Thr Thr Tyr Lys Leu>

867   GTT ATT AAT GGT AAA ACA TTG AAA GGC GAA ACA ACT ACT GAA GGT
273   Val Ile Asn Gly Lys Thr Leu Lys Gly Glu Thr Thr Thr Glu Ala>

912   GTT GAT GGT GGT ACT GCA GAA AAA GTG TTG AAA CAA TAC GGT AAC
288   Val Asp Ala Ala Thr Ala Glu Lys Val Phe Glu Gln Thr Ala Asp>

957   GAC AAC GGT GTT GAC GGT GAA TCG ACT TAC GAC GAT GCG ACT AAC
303   Asp Asn Gly Val Asp Gly Glu Thr Thr Tyr Asp Asp Ala Thr Lys>

992   ATG TTT TTA GTT TTT GAA AAA CCA GAA TTG TTT GAT GCG TTT TAA
318   Met Phe Thr Val Thr Glu Lys PTT Glu Val Ile Asp Ala Ser Ile

1047  TTA ACA CCA GCG GTG ACA AGA TCG GGT GAT AAC AAT TTG AAC AAA
133   Leu Thr Pro Ala Val Thr Arg Ser Ala Asp Asn Asn Phe Asn Lys>

1092  GAA CAA CAA AAT GGT TTG TAT GAA ATG TTG AAT ATG GGT AAC TTA
148   Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu>

1127  TAC GAA CAA CAA GCG AAT TGT TTT TTT TTT GAA AGC TTA AAA GAT GAC
163   Asn Glu Glu Gln Arg Asn Gly Pro Ile Gln Ser Leu Lys Asp Asp>

1182  TCA AGC CAA AGT GGT AAC GTA TTG TCA GAA GGT AAA AAG TTA AAT
178   Pro Ser Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn>

1227  GAA TCT CAA GCA CCG AAA GAT CCA TCC GCG TGATCAATGCACGACACA
193   Glu Ser Gln Ala Pro Lys Asp Arg Ser Ala End<

1276  TGATGATCTGCTGCTGCACTTAATTACTATGTTCTATACAAATAAATACACCGCGCT

1335  ACCGCGGTGTTCCTTATATGGTGTAAATGTAGCGAGTAATTTTAACTACTTGTGCT

1394  TCCGAAATTCACCTGCGCGGATGCTATATA

```

Table 13

Predicted nucleotide and deduced amino acid sequence for  
RDT-G-G-A.

		TCCTCTAGATCGCGCTAATTAACTAAGCTTAAAGGAGGAAAAAATT ATG AAA Met Lys>
5		
10	57 3	AGA ATA GTG CCA AAG TTC ACT GAA ATC TTC CCC GTG GAG GAC GCG Arg Ile Val Pro Lys Phe Thr Glu Ile Phe Pro Val Glu Asp Ala>
	102 18	AAC TAC CTT TAC AGC GGC TTC ATC GCG TCG GTC CCG AAA GAC GTG Asn Tyr Pro Tyr Ser Ala Phe Ile Ala Ser Val Arg Lys Asp Val>
15	147 33	ATC AAA CAC TGC ACC GAC CAT AAA GCG ATC TTC CAG CCC GTG GTG Ile Lys His Cys Thr Asp His Lys Gly Ile Phe Gln Pro Val Leu>
	192 48	CCA CCG GAG AAG AAG GTC CCG GAG CTA TGG TTC TAC ACA GAG CTC Pro Pro Glu Lys Lys Val Pro Glu Leu Trp Phe Tyr Thr Glu Leu>
	237 63	AAA ACT AGG ACC AGC TCC ATC ACC CTC CCC ATA CCG ATG GAC AAC Lys Thr Arg Thr Ser Ser Ile Thr Leu Ala Ile Arg Met Asp Asn>
20	282 78	CTG TAC CTC GTG GCG TTC AGG ACC CCG GCG GCG GTG TGG TGG GAG Leu Tyr Leu Val Gly Phe Arg Thr Pro Gly Gly Val Trp Trp Glu>
	327 93	TTC GCG AAG GAC GCG GAC ACC CAC CTC CTC GCG GAC AAC CCC AGG Phe Gly Lys Asp Gly Asp Thr His Leu Leu Gly Asp Asn Pro Arg>
25	372 108	TGG CTC GCG TTC GCG GCG AGG TAC CAG GAC CTC ATC GCG AAC AAG Trp Leu Gly Phe Gly Gly Arg Tyr Gln Asp Leu Ile Gly Asn Lys>
	417 123	GCT CTC GAG ACC GTC ACC ATG GCG CCG GCG GAA ATG ACC AGG GCG Gly Leu Glu Thr Val Thr Met Gly Arg Ala Glu Met Thr Arg Ala>
	462 138	GTC AAC CAC CTG GCG AAG AAG AAG AAG GCG GCT CAC CCA CAG GCC Val Asn Asp Leu Ala Lys Lys Lys Lys Ala Ala Asp Pro Gln Ala>
30	507 153	GAC ACC AAG ACC AAG CTG GTG AAG CTG GTG GTC ATG GTG TCC GAG Asp Thr Lys Ser Lys Leu Val Lys Leu Val Val Met Val Cys Glu>
	552 168	TGG CTC GCG TTC AAC ACC GTC TCC CCG ACC GTG GAC GCG GCG TTC Gly Leu Arg Phe Asn Thr Val Ser Arg Thr Val Asp Ala Gly Phe>
	597 183	AAC AGC CAG CAC GCG GTG ACC TTC ACC GTG ACC CAG GCG AAG CAG Asn Ser Gln His Gly Val Thr Leu Thr Val Thr Gln Gly Lys Gln>
35	642 199	CTG CAG AAG TGG GAC AGG ATC TCC AAG GCG GCG TTC GAG TGG GGT Val Gln Lys Trp Asp Arg Gln Ser Lys Ala Ala Phe Glu Trp Ala>
	687 213	GAC CAG GCG ACC GGT GTG ATC GCG CAC ATG CAG AAG GTT GCG ATC Asp His Pro Thr Ala Val Ile Pro Asp Met Gln Lys Leu Gly Ile>
40	712 219	AAG GAT AAG AAC GAA GCA GCG AGG ATC GTT GCG CTC GTT AAG AAT Lys Asp Lys Asn Glu Ala Ala Arg Ile Val Ala Leu Val Lys Asn>
	777 243	GAA ACT ACT GCG GGT GCG GGT ACT GGT GGA TCC AAA CCA GAA GTG Glu Thr Thr Ala Ala Ala Ala Thr Ala Gly Ser Lys Pro Gln Val>
	822 258	ATC GAT TGG TGT GAA TTA ACA GCA GCG GTG ACA ACT TAC AAA GTT Ile Asp Ala Ser Glu Leu Thr Pro Ala Val Thr Thr Tyr Lys Leu>
45	867 273	CTT ATT CAG GGT AAA ACA TTT AAA GCG GAA ACA ACT ACT GAA GGT Val Ile Asn Gly Lys Thr Leu Lys Gly Glu Thr Thr Thr Gln Ala>
	912 288	CTT GAT GCG GGT ACT GCA GAA AAA GTC TTC AAA CAA TAC GGT AAC Val Asp Ala Ala Thr Ala Glu Lys Val Phe Lys Gln Tyr Ala Asn>
50	957 303	GAC AAG GGT GTT CAC GGT GAA TGG ACT TAC GAG GAT GCG ACT AAG Asp Asn Gly Val Asp Gly Glu Trp Thr Tyr Asp Asp Ala Thr Lys>
	1002 319	TTC TTT TCA GTT ACT GAA AAA GCA GAA GTG ATC GAT GCG TGT GAA Thr Phe Thr Val Thr Glu Lys Pro Ile Val Ile Asp Ala Ser Glu>

Table 13 (cont.)

5

10

15

20

25

30

35

```

1047   TTA ACA CCA GCC GTC ACA AGA TCC AAA CCA GAA GTC ATC GAT GCG
333   Leu Thr Pro Ala Val Thr Arg Ser Lys Pro Glu Val Ile Asp Ala>

1092   TCT GAA TTA ACA CCA GCC GTC ACA ACT TAC AAA GTT GTT ATT AAT
348   Ser Glu Leu Thr Pro Ala Val Thr Thr Tyr Lys Leu Val Ile Asn>

1137   GGT AAA ACA TTG AAA GCC GAA ACA ACT ACT GAA GCT GTT GAT GCT
363   Gly Lys Thr Leu Lys Gly Glu Thr Thr Thr Glu Ala Val Asp Ala>

1182   GGT ACT CCA GAA AAA GTC TTC AAA CAA TAC GCT AAC GAC AAC GGT
378   Ala Thr Ala Glu Lys Val Phe Lys Gln Tyr Ala Asn Asp Asn Gly>

1227   GTT GAC GGT GAA TGG ACT TAC GAC GAT GCG ACT AAG ACC TTT ACA
393   Val Asp Gly Glu Trp Thr Tyr Asp Asp Ala Thr Lys Thr Phe Thr>

1272   GTT ACT GAA AAA CCA GAA GTC ATC GAT GCG TCT GAA TTA ACA CCA
408   Val Thr Glu Lys Pro Glu Val Ile Asp Ala Ser Glu Leu Thr Pro>

1317   TGG GTG ACA AGA TCG GGT GAT AAC AAT TTC AAC AAA GAA CAA CAA
423   Val Val Thr Trp Ser Ala Asp Asn Asn Phe Asn Lys Glu Gln Glu>

1362   AAT GGT TTC TAT GAA ATC TTC AAT ATG GGT AAC TTA AAC GAA GAA
438   Asn Ala Phe Thr Glu Ile Leu Asn Met Pro Asn Leu Asn Glu Glu>

1407   GAA GCG AAT GGT TTC ATC GAA AGC TTA AAA GAT GAC CCA AGC CAA
453   Glu Arg Asn Gly Phe Ile Leu Ser Leu Lys Asp Asp Pro Ser Glu>

1452   AAT GGT AAC GTA TTC TCA GAA GGT AAA AAG TTA AAT GAA TCT GAA
468   Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Glu>

1497   TGA GCG AAA GAT GCA TCG GCG TATCAATGCAAGGACACATCATCATGCT
483   Ala Pro Lys Asp Arg Ser Ala End<

1549   GCTGCACCTTAATTACTATGTTGGTATACAAATAAATACACCCGGCGTACGGCGGTGTTCC

1608   TATATGGTGTAAATGTAGCCACTAAATTTAACTACTTTCTGTCGCGGAATTCAC

1667   TCGGCGCATGCTATATA

```

When these genes were expressed in *E.coli* using the pGEMEX system the expected chimeric proteins were produced. The RDT-G-A produced a protein of predicted molecular weight 44,576 Daltons (pI 7.2). RDT-G-G-A produced a slightly larger polypeptide predicted to be 53,955 Daltons with a predicted pI of 5.4.

Crude bacterial lysates of cells expressing RDT-G-A or RDT-G-G-A were potent inhibitors of eukaryotic protein synthesis in the rabbit reticulocyte assay described in Example 1. Papain treatment of the lysates further increases activity. Analysis of the papain treated lysates indicates that the intact RDT domain is released from the ABR domains.

Both RDT-G-A and RDT-G-G-A bind tightly to IgG Sepharose (Pharmacia LKB Biotechnology). Binding is stable at pH 5.0. Elution was accomplished with 0.5 M ammonium acetate pH 3.5 or by boiling the resin in SDS.

#### Example 12:

Seeds from the following species of *Panicoideae* were ground to a fine consistency in a mortar and pestle (after removal of the glume if necessary): *Zea mays mays*, *Z.m. mexicana*, *Z.m. parviglumis*, *Z. luxurians*, *Z. mexicana*, *Z. mexicana*, *Tripsacum dactyloides*, *Coix lachryma-jobi*, *Sorghum bicolor*.

Soluble proteins were extracted from mature dry seed by the following techniques. The proteins were extracted for 2 hours with 50 mM sodium phosphate buffer, pH 7.5 containing 25 µg/ml leupeptin, 25 µg/ml antipain, 2 mM EDTA and 4 mM phenylmethane sulfonyl fluoride. The extraction buffer per gram seed tissue was used. After centrifugation to remove insoluble material, an aliquot of the extract was analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis using 17-27% gradient gels, then electroblotted onto PVDF paper (Millipore). The blot was probed using rabbit antisera against maize RIP α

and  $\beta$  fragments (1:2000 dilution each) as primary antibody and goat anti-rabbit IgG antibody as secondary antibody, then developed with NBT/BCIP.

Except the *Coix* extract, all the extracts tested showed cross-reactivity with the maize RIP antisera. A prominent band at ~34 kD was observed corresponding to proRIP, and bands at ~16.5 and 8.5 kD were also observed, corresponding to  $\alpha$  and  $\beta$  fragments respectively. This indicates that RIP forms equivalent to maize proRIP and the activated  $\alpha\beta$  form are present in many members of the subfamily Panicoideae.

#### Example 13:

Total DNA from the following species was isolated according to the procedures described by Saghai-Maroo et al., *PNAS*, 81:8014-8018, 1984. The following species of *Panicoideae* were included: 3 accessions of *Zea mays* ssp.; *parviglumis*; *Zea luxurians*; *Zea mays* ssp. *mexicana*; *Coix lachryma-jobi*; *Sorghum bicolor*; and *Zea mays* ssp. *mays* var. B73.

Generally, about 8  $\mu$ g of the extracted DNA from each sample was digested to completion with 20 units of *Hind* III, *Eco* RI and *Sst* I in 20 microliters of a reaction mixture containing the appropriate reaction buffer at 37 °C for 2 hours.

Next, the total extracted DNA from each sample was subjected to the Southern hybridization technique (see Southern (1975), *J.Mol.Biol.*, 98:503-517). The DNA fragments were fractionated on the basis of their size by means of electrophoresis on a 0.8% agarose gel. The double-stranded DNA fragments were modified into single-stranded DNA fragments in an alkali solution; and then a nylon filter was placed into close contact with the gel to transfer the modified DNA segments onto the filter in the presence of a high salt concentration solution.

Hybridization was carried out using, as the probe, the cDNA clone of the RIP gene (the probe fragment is provided from position 2 to 1075 in Table 1). The probe was radiolabeled with  $^{32}$ P and the signals in Southern transfers were visualized by autoradiography.

The Southern blots were hybridized with the cDNA clone of the RIP gene. A single fragment with strong homology to the clone was observed for each enzyme/species combination except the *Coix* accession. The inbred line of maize has a single major band with two minor bands. The other species which, with the exception of *Sorghum*, are not inbred showed between 2 and 5 minor bands. The *Coix* had either 4 or 5 such bands depending on the enzyme used.

Although the invention has been described in considerable detail, with reference to certain preferred embodiments thereof, it will be understood that variations and modifications can be affected within the spirit and scope of the invention as described above and as defined in the appended claims.

#### Claims

1. A substantially pure protein, termed a proRIP, wherein the protein is incapable of substantially inactivating eukaryotic ribosomes, but which can be converted into a protein, termed an RIP, that is capable of substantially inactivating eukaryotic ribosomes, said proRIP having a removable, internal peptide linker sequence.
2. The substantially pure protein of Claim 1, wherein the protein is derived from a *Panicoideae*.
3. The substantially pure protein of Claim 1, wherein the proRIP has a removable, internal peptide linker selectively inserted in the amino acid sequence of a protein capable of substantially inactivating eukaryotic ribosomes, termed an  $\alpha\beta$  RIP derived from the group consisting of *Panicoideae* RIP, barley RIP, ricin A-chain RIP, saporin RIP, abrin A-chain RIP, SLT-1 RIP,  $\alpha$ -trichosanthin RIP, Luffin-A RIP, and Mirabilis antiviral protein RIP.
4. The substantially pure protein of Claim 3, wherein the internal peptide linker has a sequence effectively homologous to the following amino acid sequence:

**MATLEEEVVKMQMQMPEAADLAAAA.**

5. The substantially pure protein of Claim 3, wherein the maize proRIP, has an amino acid sequence effectively homologous to the following sequence:

KRIVPKFTEIFPVEDANYPYSAFIASVRKDV IKHCTDHKGIFQPVLPPEKKVPELW  
 FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
 5 GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
 AVNDLAKKKKMATLEEEVKMOMOMPEAADLAAAAAADPQADTKSKLVK  
 LVVMVCEGLRFNTVSRTVDAGFNSQHGVTTLTVTQGKQVQKWDRIS  
 10 KAAFEWADHPTAVIPDMQKLGKDKNEAARIVALVKNQTTAAAATAAS  
 ADNDDDEA.

- 15 6. A substantially pure protein, termed an RIP, having  $\alpha$  and  $\beta$  fragments and being capable of substantially inactivating eukaryotic ribosomes derived from a *Panicoideae*.
7. The substantially pure protein of Claim 6, wherein the  $\alpha$  fragment has an amino acid sequence effectively homologous to the following sequence:

20 KRIVPKFTEIFPVEDANYPYSAFIASVRKDV IKHCTDHKGIFQPVLPPEKKVPELW  
 FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
 GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
 25 AVNDLAKKKK

and the  $\beta$  fragment has an amino acid sequence effectively homologous to the following sequence:

30 DPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHGVTTLTVTQGKQVQKWDRIS  
 SKAAFEWADHPTAVIPDMQKLGKDKNEAARIVAL  
 35 VKNQTTAAAATAASADNDDDEA.

8. A substantially pure protein having  $\alpha$  and  $\beta$  fragments and being capable of substantially inactivating eukaryotic ribosomes, termed an  $\alpha\beta$  RIP, wherein the  $\alpha\beta$  RIP is derived from the group consisting of *Panicoideae* RIP, barley RIP, ricin A-chain RIP, saporin RIP, abrin A-chain RIP, SLT-1 RIP, and  $\alpha$ -trichosanthin RIP, Luffin-A RIP, and Mirabilis antiviral protein RIP..
9. A fusion protein capable of substantially inactivating eukaryotic ribosomes, said protein having an amino acid sequence effectively homologous to the following sequence:

45 KRIVPKFTEIFPVEDANYPYSAFIASVRKDV IKHCTDHKGIFQPVLPPEKKVPELW  
 FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
 GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
 50 AVNDLAKKKKDPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHG  
 VTLTVTQGKQVQKWDRISKAAFEWADHPTAVIPDMQKLGKDKN  
 EAARIVALVKNQTTAAAATAASADNDDDEA.

10. A conjugate comprising a targeting vehicle and a protein, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic

ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes.

11. The conjugate of Claim 10 wherein the protein is derived from the group consisting of a *Panicoideae* RIP, barley RIP, ricin A-chain RIP, saporin RIP, abrin A-chain RIP, SLT-1 RIP, and  $\alpha$ -trichosanthin RIP, Luffin-A RIP, and Mirabilis antiviral protein RIP.

12. A conjugate comprising a targeting vehicle and a protein, termed an  $\alpha\beta$  RIP, that is capable of substantially inactivating eukaryotic ribosomes and that contains  $\alpha$  and  $\beta$  fragments.

13. A conjugate comprising a targeting vehicle and a protein derived from maize, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an RIP, that is capable of substantially inactivating eukaryotic ribosomes, said  $\alpha$  fragment having an amino acid sequence effectively homologous to the following sequence:

KRIVPKFTEIFPVEDANYPYSAFIASVRKDVIKHCTDHKGIFQPVLPPEKKVPELW  
FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
AVNDLAKKKK

and the  $\beta$  fragment having an amino acid sequence effectively homologous to the following sequence:

DPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHGVTLTVTQGKQVQKWDR  
SKAAFEWADHPTAVIPDMQKLGKDKNEAARIVAL  
VKNQTTAAAATAASADNDDDEA.

14. A conjugate comprising a targeting vehicle and a fusion protein capable of substantially inactivating eukaryotic ribosomes, said protein having an amino acid sequence effectively homologous to the following sequence:

KRIVPKFTEIFPVEDANYPYSAFIASVRKDVIKHCTDHKGIFQPVLPPEKKVPELW  
FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
GDTHLLGDNPRWLGFGGRYQDLIGNKGLETVTMGRAEMTR  
AVNDLAKKKKDPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHG

VTLTQGKQVQKWDRISKAAFEWADHPTAVIPDMQKLGKDKN  
EAARIVALVKNQTTAAAATAASADNDDDEA.

15. The conjugate of any one of Claims 10 to 15, wherein the targeting vehicle is selected from the group consisting of hormones, antibodies, antibody binding proteins, growth factors and lectins, or fragments thereof.

16. A method for converting a proRIP into an RIP, said method comprising the following steps:  
a) providing a homogeneous protein derived from the group consisting of consisting of a *Panicoideae* RIP, barley RIP, ricin A-chain RIP, saporin RIP, abrin A-chain RIP, SLT-1 RIP, and  $\alpha$ -trichosanthin RIP, Luffin-A RIP, and Mirabilis antiviral protein RIP, termed a proRIP, wherein the

proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an RIP, that is capable of substantially inactivating eukaryotic ribosomes; and

- 5 b) contacting the proRIP with a cleaving agent capable of deleting the linker to form a protein having  $\alpha$  and  $\beta$  fragments, termed an RIP, that is capable of substantially inactivating eukaryotic ribosomes.

17. A DNA isolate capable of encoding a protein, termed a proRIP, wherein the proRIP has a removable, internal peptide linker sequence and is incapable of substantially inactivating eukaryotic ribosomes, but  
10 which upon removal of the linker can be converted into a protein having  $\alpha$  and  $\beta$  fragments, termed an RIP, that is capable of substantially inactivating eukaryotic ribosomes.

18. The DNA isolate of Claim 17, wherein the DNA isolate encodes a protein derived from the group consisting of consisting of *Panicoideae* RIP, barley RIP, ricin A-chain RIP, saporin RIP, abrin A-chain  
15 RIP, SLT-1 RIP, and  $\alpha$ -trichosanthin RIP, Luffin-A RIP, and Mirabilis antiviral protein RIP.

19. The DNA isolate of Claim 17, wherein the DNA isolate encodes the maize proRIP of Claim 5.

20. A DNA isolate encoding a fused protein capable of substantially inactivating eukaryotic ribosomes, said  
20 protein having an amino acid sequence effectively homologous to the following sequence:

KRIVPKFTEIFFVEDANYPYSAFIASVRKDVIKHCTDHKGIFQPVLPPEKKVPELW  
FYTELKTRTSSITLAIRMDNLYLVGFRTPGGVWWEFGKD  
25 GDTHLLGDNPRWLGGGGRYQDLIGNKGLETVTMGRAEMTR  
AVNDLAKKKKDPQADTKSKLVKLVVMVCEGLRFNTVSRTVDAGFNSQHG  
VTLTVTQGKQVQKWDRISKAAFEWADHPTAVIPDMQKLGKIDKN  
30 EAARIVALVKNQTTAAAATAASADNDDDEA.

21. A DNA isolate encoding an RIP having at least one sites engineered therein, whereby the DNA isolate  
35 may be cleaved by restriction enzymes to permit insertion of a DNA sequence encoding a linker.

22. A biologically functional expression vehicle containing a DNA isolate of any one of Claims 17 to 21.

23. A host cell transformed with a biologically functional expression vehicle of Claim 22.


- 40 24. The transformed host cell of Claim 23, wherein the host cell is a eukaryotic cell.

25. The host cell of Claim 24, wherein the host cell is maize.

26. A method of creating a protein incapable of substantially inactivating eukaryotic ribosomes, termed a  
45 proRIP, said method comprising the steps of


- (a) providing a first DNA sequence encoding an RIP having at least one sites engineered therein,  
(b) cleaving the first DNA sequence with a restriction enzyme to form first DNA subsequences,  
(c) providing a second DNA isolate encoding a polypeptide of desired length and amino acid residues  
50 and having ends ligatable with the cleaved ends of the first DNA subsequences,  
(d) ligating the first DNA subsequences and the second DNA to form a third DNA sequence  
capable of expressing a proRIP,  
(e) expressing the proRIP.

**FIG. 1** *Processing of maize proRIP to active form*





**FIG. 2** Inhibition of protein synthesis in a rabbit reticulocyte lysate by purified maize RIP.





European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 91201436.2
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.5)
Y	GB - A - 2 216 891 (FARMITALIA CARLO ERBA S.r.l.) * Example; claims *	1,10, 15,17, 22,23	C 07 K 15/10 C 12 N 15/29 C 12 N 5/10
D,Y	EUROPEAN JOURNAL OF BIOCHE- MISTRY, vol. 148, 1985, New York, Berlin F. I. LAMB et al. " Nucleotide sequence of cloned cDNA coding for preproricin" pages 265-270 * Discussion *	1,3, 16,17	
D,A	CANCER SURVEYS, vol. 1, no. 3, 1982, Oxford, England L. BARBIERI and F. STIRPE "Ribosome-inactivating pro- teins from plants: Properties and possible uses" pages 489-520 * Page 511-512 *	1,22- 26	
			TECHNICAL FIELDS SEARCHED (Int. CL.5)
			C 07 K C 12 N A 61 K
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 23-08-1991	Examiner AUGUSTIN
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	